

Short notes for the course of Functional Analysis, PhD in Statistics

Annalisa Cesaroni

Contents

1	Introduction	1
2	Measure theory and integration	3
2.1	Measure space	3
2.2	Borel measures on \mathbb{R} , and the Lebesgue measures on \mathbb{R} and \mathbb{R}^n	4
2.3	Some interesting examples	7
2.4	Measurable functions	7
2.5	Integration with respect to the Lebesgue measure and L^p spaces	8
2.6	Decomposition of measures	9
2.7	Distribution of a random variable	10
2.8	Problems	11
3	Banach spaces	13
3.1	Banach spaces	13
3.2	Bounded linear operators	14
3.3	L^p spaces	15
3.4	Spaces of random variables with finite moments	17
3.5	Modes of convergence and compactness theorems	18
3.6	Problems	20
4	Hilbert spaces	22
4.1	Hilbert spaces	22
4.2	Orthogonality and projections in Hilbert spaces	22
4.3	Hilbert space of random variables	24
4.4	The Hilbert spaces L^2 and H^1	25
4.5	Bounded linear operators in Hilbert spaces	26
4.6	Problems	29
	References	30
A	Solutions to problems Section 2	30
B	Solutions to problems Section 3	32
C	Solutions to problems Section 4	36

1 Introduction

These notes are intended for the first year students of the PhD course in Statistics, at University of Padova. They are not exhaustive, nor complete, but they could serve as a basis of the study of the arguments presented during the course of Functional Analysis. The topics are presented in a

quite informal way, trying to reach also students without a specific preparation in mathematics. Only few proofs are provided and for the others bibliographical references are provided. The notes are divided in 2 parts: the first part contains a review of results of Real Analysis, in particular on measure theory and integration with respect to the Lebesgue measure, whereas the second part is a brief review of basic results in linear Functional Analysis, in particular on Banach and Hilbert spaces, and to the main examples, the L^p spaces and the L^2 space. At the end of each section some exercises are proposed, more or less simple to solve. In the appendix there are the (sketchy) solutions to the problems.

2 Measure theory and integration

2.1 Measure space

We fix a set X and we define $\mathcal{P}(X)$ the set of all subsets of X .

Definition 2.1. $\Sigma \subset \mathcal{P}(X)$ is a σ -algebra on X if

- it is closed by complement, that is if $A \in \Sigma$ then $X \setminus A \in \Sigma$,
- it is closed by countable union, that is if $(A_i)_i$ is a sequence of elements in Σ then $\cup_{i=1}^{\infty} A_i \in \Sigma$.

Let $\mathcal{C} \subseteq \mathcal{P}(X)$, then $\Sigma(\mathcal{C})$, the σ -algebra generated by \mathcal{C} is the smallest σ -algebra which contains all the elements in \mathcal{C} (and then all countable intersections and countable unions of elements in \mathcal{C}).

The smallest possible σ -algebra on X is given by $\Sigma = \{\emptyset, X\}$, and the largest possible σ -algebra on X is $\Sigma = \mathcal{P}(X)$.

Definition 2.2. $\mathcal{B}(\mathbb{R})$ is the σ -algebra on \mathbb{R} generated by all the intervals $\mathcal{C} = \{(a, b) \mid a, b \in \mathbb{R}\}$. $\mathcal{B}(\mathbb{R}^N)$ is the σ -algebra on \mathbb{R}^N generated by all the pluri-rectangulars $\mathcal{C} = \{\prod_{i=1}^N (a_i, b_i) \mid a_i, b_i \in \mathbb{R}\}$.

Remark 2.3. Note that $\sigma(\mathcal{C}) = \mathcal{B}(\mathbb{R})$ also when $\mathcal{C} = \{(a, b] \mid a, b \in \mathbb{R}\}$, since $(a, b) = \cup_n (a, b - \frac{1}{n}]$, or when $\mathcal{C} = \{[a, b) \mid a, b \in \mathbb{R}\}$, since $(a, b) = \cup_n [a + \frac{1}{n}, b)$, or when $\mathcal{C} = \{[a, b] \mid a, b \in \mathbb{R}\}$ again because $(a, b) = \cup_n [a + \frac{1}{n}, b - \frac{1}{n}]$. Analogously $\sigma(\mathcal{C}) = \mathcal{B}(\mathbb{R})$ when $\mathcal{C} = \{(a, +\infty) \mid a \in \mathbb{R}\}$, since $(a, b] = (a, +\infty) \cap (-\infty, b]$, and $(-\infty, b] = \mathbb{R} \setminus (b, +\infty)$ and so on.

Definition 2.4. Let Σ be a σ -algebra on X . A function $\mu : \Sigma \rightarrow [0, +\infty]$ is a measure if

- $\mu(\emptyset) = 0$,
- it is σ -additive, that is if $(A_i)_i$ is a sequence of elements in Σ with $A_i \cap A_j = \emptyset$ for $i \neq j$ then $\mu(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mu(A_i)$.

(X, Σ, μ) is called a measure space.

If $\mu(X) < +\infty$, then μ is a finite measure (a probability measure if $\mu(X) = 1$). Usually measure spaces with probability measures are denoted with Ω (in place of X), the σ -algebra is \mathcal{F} (in place of Σ) and the measure is \mathbb{P} (in place of μ).

If $X = \cup_i A_i$, with $\mu(A_i) < +\infty$ for all i , μ is σ -finite.

If $X = \mathbb{R}^n$, $n \geq 1$ and $\Sigma = \mathcal{B}(\mathbb{R}^n)$, then μ is called a Borel measure.

Example 2.5. Let $x_0 \in \mathbb{R}$, and define the measure on $\mathcal{P}(\mathbb{R})$ as $\delta_{x_0}(A) = \begin{cases} 1 & x_0 \in A \\ 0 & x_0 \notin A \end{cases}$.

Then δ_{x_0} is called Dirac measure centered at x_0 .

Proposition 2.6 (Monotonicity, subadditivity, continuity). Let μ be a measure on Σ . Then

- (i) if $A \subset B$, $A, B \in \Sigma$, then $\mu(A) \leq \mu(B)$ (monotonicity with respect to inclusion);
- (ii) if $(A_i)_i$ is a sequence of elements in Σ then $\mu(\cup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \mu(A_i)$;
- (iii) if $(A_i)_i$ is a sequence of elements in Σ with $A_i \subseteq A_{i+1}$ then $\mu(\cup_{i=1}^{\infty} A_i) = \lim_{i \rightarrow +\infty} \mu(A_i)$;
- (iv) if $(A_i)_i$ is a sequence of elements in Σ with $A_i \supseteq A_{i+1}$ and $\mu(A_{i_0}) < +\infty$ for some i_0 , then $\mu(\cap_{i=1}^{\infty} A_i) = \lim_{i \rightarrow +\infty} \mu(A_i)$.

Proof. (i) Observe that $B = A \cup (B \setminus A)$, so by σ -additivity $\mu(B) = \mu(A) + \mu(B \setminus A) \geq \mu(A)$.

(ii) Let $B_1 = A_1$ and $B_i = A_i \setminus \cup_{k=1}^{i-1} A_k$ then B_i are disjoint and

$$\mu(\cup_i A_i) = \mu(\cup_i B_i) = \sum_{i=1}^{+\infty} \mu(B_i) \leq \sum_{i=1}^{+\infty} \mu(A_i).$$

(iii) Let $B_1 = a_1$ and $B_i = A_i \setminus A_{i-1}$ then

$$\mu(\cup_i A_i) = \mu(\cup_i B_i) = \sum_{i=1}^{+\infty} \mu(B_i) = \lim_{n \rightarrow +\infty} \sum_{i=1}^n \mu(B_i) = \mu(A_n).$$

(iv) Let $F_i = A_{i_0} \setminus A_i$ for $i > i_0$. Then $\mu(A_{i_0}) = \mu(F_i) + \mu(A_i)$, $F_i \subseteq F_{i+1}$ and $\cup_i F_i = A_{i_0} \setminus \cap_i A_i$. Therefore by 1), we get

$$\mu(A_{i_0}) = \mu(\cap_i A_i) + \lim_i \mu(F_i) = \mu(\cap_i A_i) + \lim_i (\mu(A_{i_0}) - \mu(A_i))$$

and we cancel $\mu(A_{i_0})$ from both sides. □

Definition 2.7. Let (X, Σ, μ) a measure space. The completion of Σ with respect to μ is the σ -algebra

$$\mathcal{M} = \{A \subseteq X \mid \exists B, C \in \Sigma, \mu(C) = 0, B \subseteq A, A \setminus B \subseteq C\}.$$

Definition 2.8. Let (X, Σ, μ) a measure space. A property holds **almost everywhere** if there exists $N \in \Sigma$ with $\mu(N) = 0$ such that the property holds for all $x \in X \setminus N$.

Proposition 2.9. Let Σ be a σ -algebra on X and $\mu : \Sigma \rightarrow [0, +\infty]$ with $\mu(\emptyset) = 0$. Then they are equivalent:

- (i) μ is σ -additive: if $(A_i)_i$ is a sequence of elements in Σ with $A_i \cap A_j = \emptyset$ for $i \neq j$ then $\mu(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mu(A_i)$,
- (ii) μ is additive: if $A, B \in \Sigma$ and $A \cap B = \emptyset$ then $\mu(A \cup B) = \mu(A) + \mu(B)$
and
 μ is countable subadditive: if $(A_i)_i$ is a sequence of elements in Σ then $\mu(\cup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \mu(A_i)$;
- (iii) μ is additive: if $A, B \in \Sigma$ and $A \cap B = \emptyset$ then $\mu(A \cup B) = \mu(A) + \mu(B)$
and
 μ is continuous on increasing sequence of sets: if $(A_i)_i$ is a sequence of elements in Σ with $A_i \subseteq A_{i+1}$ then $\mu(\cup_{i=1}^{\infty} A_i) = \lim_{i \rightarrow +\infty} \mu(A_i)$.

Proof. The fact that (i) implies (ii) and that (i) implies (iii) has been proved in Proposition 2.6. We prove that (ii) implies (i). We consider a sequence $(A_i)_i$ of elements in Σ with $A_i \cap A_j = \emptyset$ for $i \neq j$. Then by (ii) we get that $\mu(\cup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \mu(A_i)$. On the other hand by additivity and monotonicity (which is a consequence of additivity) we get that for every n , $\mu(\cup_{i=1}^{\infty} A_i) \geq \mu(\cup_{i=1}^n A_i) = \sum_{i=1}^n \mu(A_i)$. Sending $n \rightarrow +\infty$ we conclude $\mu(\cup_{i=1}^{\infty} A_i) \geq \sum_{i=1}^{\infty} \mu(A_i)$.

We prove that (iii) implies (i). We consider a sequence $(A_i)_i$ of elements in Σ with $A_i \cap A_j = \emptyset$ for $i \neq j$. We define $B_i = \cup_{j=1}^i A_j$. Then $\cup_i B_i = \cup_i A_i$. Note that by additivity $\mu(B_i) = \sum_{j=1}^i \mu(A_j)$ and that $B_1 \subseteq B_2 \subseteq B_3 \dots$. Therefore by (iii) and additivity we get

$$\mu(\cup_{i=1}^{\infty} A_i) = \mu(\cup_{i=1}^{\infty} B_i) = \lim_{i \rightarrow +\infty} \mu(B_i) = \lim_{i \rightarrow +\infty} \sum_{j=1}^i \mu(A_j) = \sum_{j=1}^{\infty} \mu(A_j).$$

□

2.2 Borel measures on \mathbb{R} , and the Lebesgue measures on \mathbb{R} and \mathbb{R}^n .

Let $F : \mathbb{R} \rightarrow \mathbb{R}$ be an increasing function which is right continuous, that is $\lim_{x \rightarrow a^+} F(x) = F(a)$. We define for all $a, b \in \mathbb{R}$,

$$\mu_F(a, b] = F(b) - F(a).$$

Moreover $\mu_F(\emptyset) = 0$ and for all sequences $a_1 < b_1 < a_2 < b_2 < \dots < a_i < b_i < a_{i+1} < b_{i+1} \dots$

$$\mu_F(\cup_i(a_i, b_i]) = \sum_i F(b_i) - F(a_i).$$

Observe that if we define $\mathcal{C} = \{(a, b], a, b \in \mathbb{R}\}$, then $\Sigma(\mathcal{C}) = \mathcal{B}(\mathbb{R})$. Note that if $F_1 = F_2 + c$ for some constant then $\mu_{F_1} = \mu_{F_2}$. Also the viceversa is true: if $\mu_{F_1} = \mu_{F_2}$, then $F_1 = F_2 + c$ for some constant c .

Remark 2.10. Note that F monotone increasing implies that $\mu_F(a, b] \geq 0$, and moreover, since F is right continuous, then

$$\mu_F(\cup_n(a + 1/n, b]) = \mu_F(a, b] = F(b) - F(a) = F(b) - \lim_n F(a + 1/n) = \lim_n \mu_F(a + 1/n, b].$$

Reasoning as before, it is possible to see that, at least when restricted to \mathcal{C} , there holds that μ_F has positive values, is additive and is continuous with respect to increasing sequences of sets (which is enough to get σ -additivity if μ_F is defined on a σ -algebra, see Proposition 2.9).

We recall that F is monotone increasing and then $\lim_{x \rightarrow +\infty} F(x) = \sup F$ and $\lim_{x \rightarrow -\infty} F(x) = \inf F$ (we say that if $F(\mathbb{R})$ is unbounded from above, $\sup F = +\infty$ and if $F(\mathbb{R})$ is unbounded from below, $\inf F = -\infty$).

We may extend μ_F to intervals obtained by unions and intersections of elements in \mathcal{C} , and using additivity and continuity. In particular we get

$$\begin{aligned} \mu_F(a, +\infty) &= \mu_F(\cup_n(a, a + n]) = \lim_n F(a + n) - F(a) = \sup F - F(a) \\ \mu_F(-\infty, b] &= \mu_F(\cup_n(b - n, b]) = \lim_n F(b) - F(b - n) = F(b) - \inf F \\ \mu_F(a, b) &= \mu_F(\cup_{n \geq n_0}(a, b - 1/n]) = \lim_n F(b - 1/n) - F(a) = \lim_{x \rightarrow b^-} F(x) - F(a) \\ \mu_F(-\infty, b) &= \mu_F((-\infty, b - 1] \cup (b - 1, b)) = \mu_F((-\infty, b - 1]) + \mu_F((b - 1, b)) \\ &= \lim_{x \rightarrow b^-} F(x) - F(b - 1) + F(b - 1) - \inf F = \lim_{x \rightarrow b^-} F(x) - \inf F \\ \mu_F[a, b) &= \mu_F[(a - 1, b) \setminus (a - 1, a)] = \mu_F(a - 1, b) - \mu_F(a - 1, a) \\ &= \lim_{x \rightarrow b^-} F(x) - F(a - 1) - \lim_{x \rightarrow a^-} F(x) + F(a - 1) = \lim_{x \rightarrow b^-} F(x) - \lim_{x \rightarrow a^-} F(x) \\ \mu_F[a, b] &= \mu_F([a, b + 1) \setminus (b, b + 1)) = \mu_F[a, b + 1) - \mu_F(b, b + 1) \\ &= F(b) - \lim_{x \rightarrow a^-} F(x) \\ \mu_F[a, +\infty) &= \sup_F - \lim_{x \rightarrow a^-} F(x). \end{aligned}$$

In particular we note that

$$\begin{aligned} \mu_F(\mathbb{R}) &= \mu_F(\cup_n(a - n, b + n]) = \lim_n F(b + n) - F(a - n) = \sup F - \inf F \\ \mu_F(\{a\}) &= \mu_F((c, a] \setminus (c, a)) \\ &= \mu_F((c, a]) - \mu_F((c, a)) = F(a) - F(c) - (\lim_{x \rightarrow a^-} F(x) - F(c)) = F(a) - \lim_{x \rightarrow a^-} F(x). \end{aligned}$$

Theorem 2.11. (i) *There exists a unique Borel measure $\overline{\mu}_F$ which coincides with μ_F on intervals $(a, b]$. This measure is σ -finite and it is finite if and only if $\sup F - \inf F < +\infty$.*

(ii) *Given a Borel measure on \mathbb{R} which is σ -finite, there exists F monotone increasing and right continuous such that $\mu = \mu_F$.*

Proof. (i) The proof is based on the Caratheodory criterion, and we refer to [2, Theorem 1.14, Theorem 1.16]. As for the σ -finiteness it is sufficient to observe that $\nu_F(-n, n] = F(n) - F(-n) < +\infty$ and $\mathbb{R} = \cup_n(-n, n]$. Moreover, since $\mu_F(\mathbb{R}) = \sup F - \inf F$, we conclude that F is finite iff $\sup F - \inf F < +\infty$.

(ii) We want to construct F . Put $F(0) = 0$ and

$$F(x) = \begin{cases} \mu(0, x] & x > 0 \\ -\mu(x, 0] & x < 0. \end{cases}$$

Observe that if $b > a \geq 0$, $F(b) - F(a) = \mu(0, b] - \mu(0, a] = \mu(0, b] \setminus (0, a] = \mu(a, b] \geq 0$, if $0 \geq b > a$, then $F(b) - F(a) = -\mu(b, 0] + \mu(a, 0] = \mu(a, 0] \setminus (b, 0] = \mu(a, b] \geq 0$ and finally if $a < 0 < b$, then $F(b) - F(a) = \mu(0, b] + \mu(a, 0] = \mu(a, b] \geq 0$. So F is increasing.

We check that it is right continuous. First of all observe that for $a > 0$, $\lim_{x \rightarrow a^+} F(x) = \lim_n F(a + 1/n) = \lim_n \mu(0, a + 1/n] = \mu(\cap_n (0, a + 1/n]) = \mu(0, a] = F(a)$. If $a = 0$ $\lim_{x \rightarrow 0^+} F(x) = \lim_n F(1/n) = \lim_n \mu(0, 1/n] = \mu(\cap_n (0, 1/n]) = \mu(\emptyset) = 0 = F(0)$. Finally if $a < 0$, then $\lim_{x \rightarrow a^+} F(x) = \lim_n F(a + 1/n) = -\lim_n \mu(a + 1/n, 0] = -\mu(\cup_n (a + 1/n, 0]) = -\mu(a, 0] = F(a)$.

Finally we already checked that $\mu(a, b] = F(b) - F(a)$ and then we conclude that $\mu = \mu_F$.

Note that if μ is finite, we could also define (in an equivalent way) F such that $\inf F = -\infty$ and $F(x) = \mu(-\infty, x]$. □

Definition 2.12. Let $F(x) = x$ for all x , then $\bar{\mu}_F$ is called **Lebesgue measure**. We indicate with \mathcal{L} . We denote with \mathcal{M} the completion of $\mathcal{B}(\mathbb{R})$ with respect to \mathcal{L} , and we call it the Σ -algebra of Lebesgue measurable sets.

Proposition 2.13. The Lebesgue measure

- (i) associates to each interval its length,
- (ii) is translation invariant, that is $\mathcal{L}(A + x) = \mathcal{L}(A)$ for all $x \in \mathbb{R}$, $A \in \mathcal{M}$,
- (iii) is homogenous, that is $\mathcal{L}(\lambda A) = \lambda \mathcal{L}(A)$ for all $\lambda > 0$, $A \in \mathcal{M}$,
- (iv) assigns measure 0 to points, and so also to countable sets (e.g. \mathbb{Q}),
- (v) it is σ -finite, since $\mathbb{R} = \cup_{n \in \mathbb{N}} (-n, n)$ and $\mathcal{L}(-n, n) = 2n$.

Proof. The proof is immediate by definitions and σ -additivity. Exercise. □

It is possible to define the Lebesgue measures on \mathbb{R}^n as the product measure of the Lebesgue measure on \mathbb{R} . It is a Borel measure and we denote with \mathcal{M} the Σ -algebra of Lebesgue measurable sets. We refer to [2, Section 2.6].

Proposition 2.14. The Lebesgue measure on \mathbb{R}^n

- (i) associates to each set its volume,
- (ii) is translation invariant, that is $\mathcal{L}(A + x) = \mathcal{L}(A)$ for all $x \in \mathbb{R}^n$, $A \in \mathcal{M}$,
- (iii) is n -homogenous, that is $\mathcal{L}(\lambda A) = \lambda^n \mathcal{L}(A)$ for all $\lambda > 0$, $A \in \mathcal{M}$, in particular $\mathcal{L}(B(0, r)) = r^n \mathcal{L}(B(0, 1))$, where $B(0, r)$ is the ball of radius r centered at 0,
- (iv) it is σ -finite, since $\mathbb{R}^n = \cup_{k \in \mathbb{N}} B(0, k)$ and $\mathcal{L}B(0, k) = k^n \mathcal{L}(B(0, 1))$.

2.3 Some interesting examples

Measurable sets in \mathbb{R} which contain at least one interval (they are called sets with non empty interior) have positive measure. On the other hand sets which are given by countable union of isolated points have measure zero. Nevertheless there are sets with empty interior in \mathbb{R} (so that do not contain any interval) and with positive measure (almost full measure).

Example 2.15 (A set of positive measure which does not contain any interval). Let (r_n) be an enumeration of $\mathbb{Q} \cap [0, 1]$ and fix $\varepsilon > 0$ small.

Set $A = \cup_n (r_n - \varepsilon 2^{-n}, r_n + \varepsilon 2^{-n})$. Then by subadditivity, $\mathcal{L}(A) \leq \sum_n 2\varepsilon 2^{-n} = 4\varepsilon$. Moreover $B = [0, 1] \setminus A$ is a set which does not contain any interval (otherwise it should contain some rational number but $\mathbb{Q} \cap [0, 1] \subseteq A$), and moreover $\mathcal{L}(B) \geq 1 - 4\varepsilon > 0$.

Not all the subsets of \mathbb{R} are contained in $\mathcal{M}(\mathbb{R})$, so there are sets which are not measurable. This is due to the fact that if we want to define a measure μ on the intervals of \mathbb{R} such that $\mu([0, 1]) = 1$, $\mu(A \cup B) = \mu(A) + \mu(B)$ if $A \cap B = \emptyset$ and $\mu(A) = \mu(B)$ if B can be obtained translating and rotating A , then the σ -algebra of measurable sets cannot be $\mathcal{P}(\mathbb{R})$.

Example 2.16 (A set which is not (Lebesgue) measurable). We say that $x, y \in [0, 1]$ are equivalent if $x - y \in \mathbb{Q}$. Let $P \in [0, 1]$ a set such that P consists of exactly one representative point from each equivalence class (this set exists by the axiom of choice). In particular this means that if $p, p' \in P$, $p \neq p'$, then $p - p' \notin \mathbb{Q}$. We claim that P provides the required example of a non measurable set. We prove it by contradiction, showing that it is not possible for P to be measurable.

For each $q \in \mathbb{Q} \cap [0, 1]$, define

$$P_q = [(P+q) \cap [0, 1]] \cup [(P+q) \setminus [0, 1)] - 1 = \{p+q, p \in P \cap [0, 1-q)\} \cup \{p+q-1, p \in P \cap [1-q, 1)\}.$$

So P_q is obtained by considering $P + q$ and then shifting back of 1 unit the part of $P + q$ which is outside the interval $[0, 1]$.

First of all we observe that $\mathcal{L}(P) = \mathcal{L}(P_q)$. Indeed $[(P + q) \cap [0, 1]] \cap [(P + q) \setminus [0, 1)] - 1 = \emptyset$, since if $p + q \in [0, 1)$ for some $p \in P$ and $p' + q - 1 \in [0, 1)$ for some $p' \in P$, then necessarily $p + q \neq p' + q - 1$, since $p, p' \in [0, 1)$.

Moreover we observe that if $r \neq q \in \mathbb{Q} \cap [0, 1]$, then $P_r \cap P_q = \emptyset$. Indeed assume it is not true and $x \in P_r \cap P_q$, this means that $x = p + r = p' + q$, for some $p, p' \in P$ or $x = p + r = p' + q - 1$, or $x = p + r - 1 = p' + q$. In any case we get that $p - p' \in \mathbb{Q}$, which implies that $p = p'$ by definition of the set P and so $r = q$.

Finally we observe that $\cup_{q \in \mathbb{Q} \cap [0, 1]} P_q = [0, 1)$. Indeed take $x \in [0, 1)$, then there exists $p \in P$ such that x is equivalent to p , which means that there exists $q \in \mathbb{Q}$ such that $x = p + q$. In particular this implies that $q \in (0, 1]$ and $x \in P_q$.

We conclude by σ -additivity that

$$1 = \mathcal{L}([0, 1)) = \mathcal{L}(\cup_{q \in \mathbb{Q} \cap [0, 1]} P_q) = \sum_{q \in \mathbb{Q} \cap [0, 1]} \mathcal{L}(P_q) = \sum_{q \in \mathbb{Q} \cap [0, 1]} \mathcal{L}(P) = \begin{cases} 0 & \text{if } \mathcal{L}(P) = 0 \\ +\infty & \text{if } \mathcal{L}(P) > 0 \end{cases}$$

which is not possible.

2.4 Measurable functions

Definition 2.17. Let (X, Σ, μ) be a measure space, and let $f : X \rightarrow \mathbb{R}$ be a function. Then f is measurable if for all $t \in \mathbb{R}$,

$$A(t) := \{x \in X \mid f(x) > t\} = f^{-1}(t, +\infty) \in \Sigma.$$

In particular we will be interested in the case in which $(X, \Sigma, \mu) = (\mathbb{R}^n, \mathcal{M}, \mathcal{L})$. In this case saying that $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is measurable is equivalent to require that for all $A \in \mathcal{B}(\mathbb{R})$, $f^{-1}(A) \in \mathcal{M}$.

Example 2.18. Let $A \in \mathcal{M}$ and define the characteristic function of A as $\chi_A(x) = \begin{cases} 1 & x \in A \\ 0 & x \notin A. \end{cases}$ Then χ_A is measurable. Indeed $A(t) = \emptyset$ for $t \geq 1$, $A(t) = \mathbb{R}^n$ for $t \leq 0$ and $A(t) = A$ for $t \in (0, 1)$.

Example 2.19 (Random variables). If $(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space (that is a measure space endowed with a probability measure), the measurable functions, that is functions $f : \Omega \rightarrow \mathbb{R}$ such that for all $t \in \mathbb{R}$, $A(t) := \{\omega \in \Omega \mid f(\omega) > t\} \in \mathcal{F}$, are called **random variables**. Usually random variables are indicated with X instead of f .

There is a notion of convergence of measurable functions which is quite used in probability.

Definition 2.20 (Convergence in measure). Let f_n, f be measurable functions defined on the measure space (X, Σ, μ) . Then f_n converge to f in measure if for every $\varepsilon > 0$

$$\lim_n \mu\{x \in X \mid |f_n(x) - f(x)| \geq \varepsilon\} = 0.$$

If we are in a probability space, this convergence is called **convergence in probability**, since it reads

$$\lim_n \mathbb{P}\{\omega \in \Omega \mid |X_n(\omega) - X(\omega)| \geq \varepsilon\} = 0.$$

2.5 Integration with respect to the Lebesgue measure and L^p spaces

Definition 2.21. Let $k \geq 1$, A_1, \dots, A_k a finite family of disjoint sets in \mathcal{M} and $c_1, \dots, c_k > 0$. The function $\phi(x) = \sum_{i=1}^k c_i \chi_{A_i}(x)$ is called **simple function**. It is a measurable (positive) function and we define its integral as

$$\int_{\mathbb{R}^n} \phi(x) dx = \sum_{i=1}^k c_i \mathcal{L}(A_i).$$

Definition 2.22 (Lebesgue integral). Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a measurable function such that $f(x) \geq 0$ for all x . Then

$$\int_{\mathbb{R}^n} f(x) dx = \sup \left\{ \int_{\mathbb{R}^n} \phi(x) dx \mid \phi \text{ simple function with } \phi \leq f \right\}.$$

If f is not positive we define its positive part $f^+(x) = \max(f(x), 0)$ and its negative part $f^-(x) = \max(-f(x), 0)$ and we define

$$\int_{\mathbb{R}^n} f(x) dx = \int_{\mathbb{R}^n} f^+(x) dx - \int_{\mathbb{R}^n} f^-(x) dx.$$

Note that $\int_{\mathbb{R}^n} |f(x)| dx = \int_{\mathbb{R}^n} f^+(x) dx + \int_{\mathbb{R}^n} f^-(x) dx$.

Since $f^+ \leq |f|, f^- \leq |f|$, we have that

$$\left| \int_{\mathbb{R}^n} f(x) dx \right| < +\infty \quad \text{iff} \quad \int_{\mathbb{R}^n} |f(x)| dx < +\infty.$$

We denote

$$L^1(\mathbb{R}^n) := \{f : \mathbb{R}^n \rightarrow \mathbb{R} \mid f \text{ is measurable and } \int_{\mathbb{R}^n} |f(x)| dx < +\infty\}.$$

Proposition 2.23. The following properties hold.

- If $f = 0$ almost everywhere then $\int_{\mathbb{R}^n} f(x) = 0$. If $f \geq 0$ is measurable and $\int_{\mathbb{R}^n} f(x) dx = 0$ then $f = 0$ almost everywhere.

- If f, g are measurable functions such that $f = g$ almost everywhere, then $\int_{\mathbb{R}^n} f(x)dx = \int_{\mathbb{R}^n} g(x)dx$.
- If $f, g \in L^1(\mathbb{R}^n)$, $\alpha, \beta \in \mathbb{R}$, then $\int_{\mathbb{R}^n} \alpha f(x) + \beta g(x)dx = \alpha \int_{\mathbb{R}^n} f(x)dx + \beta \int_{\mathbb{R}^n} g(x)dx$.
- If $f, g \in L^1(\mathbb{R}^n)$, and $f \leq g$ then $\int_{\mathbb{R}^n} f(x)dx \leq \int_{\mathbb{R}^n} g(x)dx$.

Proof. The proof is obtained by exploiting definitions, see [2, Section 2..2] □

Remark 2.24. [On the definition of L^1] Note that due to the previous proposition, in particular the fact that if f, g are measurable functions such that $f = g$ almost everywhere, then $\int_{\mathbb{R}^n} f(x)dx = \int_{\mathbb{R}^n} g(x)dx$, **we identify functions in $L^1(\mathbb{R}^n)$ which coincide almost everywhere.** So a function f in $L^1(\mathbb{R}^n)$ is actually a class of equivalence of functions, we do not distinguish functions which are different on sets of measure zero.

Theorem 2.25 (Monotone convergence). *Let $f_k : \mathbb{R}^n \rightarrow \mathbb{R}$ measurables, positive, i.e. $f_k \geq 0$ for all k , and such that $f_k(x) \leq f_{k+1}(x)$ for all x and for all k . Then*

$$\lim_k \int_{\mathbb{R}^n} f_k(x)dx = \int_{\mathbb{R}^n} \lim_k f_k(x)dx.$$

Proof. See [2, Theorem 2.14]. □

Proposition 2.26. *An equivalent definition of the Lebesgue integral (which can be very useful) is the following. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ measurable and positive. Let for every $t > 0$ $F(t) = \mathcal{L}(A(t)) = \mathcal{L}\{x \mid f(x) > t\}$. F is called the **repartition function of f** . Then*

$$\int_{\mathbb{R}^n} f(x)dx = \int_0^{+\infty} F(t)dt.$$

Proof. See [2, Proposition 6.24] □

Definition 2.27 (L^p spaces). *We define for $p \geq 1$,*

$$L^p(\mathbb{R}^n) = \left\{ f : \mathbb{R}^n \rightarrow \mathbb{R} \mid f \text{ is measurable and } \int_{\mathbb{R}^n} |f(x)|^p dx < +\infty \right\}.$$

Note that also functions in L^p which differ on sets of measure zero are identified.

If $A \in \mathcal{M}$, then we define

$$L^p(A) = \left\{ f : \mathbb{R}^n \rightarrow \mathbb{R} \mid f \text{ is measurable and } \int_{\mathbb{R}^n} |f(x)|^p \chi_A(x) = \int_A |f(x)|^p dx < +\infty \right\}.$$

We define

$$L^\infty(\mathbb{R}^n) = \{f : \mathbb{R}^n \rightarrow \mathbb{R} \mid f \text{ is measurable and there exists } c > 0 \text{ such that } |f(x)| \leq c \text{ for almost every } x\}$$

and analogously

$$L^\infty(A) = \{f : \mathbb{R}^n \rightarrow \mathbb{R} \mid f \text{ is measurable and there exists } c > 0 \text{ such that } |f(x)| \leq c \text{ for almost every } x \in A\}.$$

2.6 Decomposition of measures

Definition 2.28. *Let ν, ρ be measures defined on $(\mathbb{R}^n, \mathcal{M})$.*

*ν is **absolutely continuous** with respect to \mathcal{L} , and we write $\nu \ll \mathcal{L}$ if $\nu(A) = 0$ for all $A \in \mathcal{M}$ such that $\mathcal{L}(A) = 0$.*

*ρ is **singular** with respect to \mathcal{L} , and we write $\rho \perp \mathcal{L}$, if there exist $A, B \in \mathcal{M}$, $A \cap B = \emptyset$, $A \cup B = \mathbb{R}^n$, such that $\mathcal{L}(A) = 0$ and $\rho(B) = 0$.*

Example 2.29. Let $x_0 \in \mathbb{R}$ and consider the Dirac measure δ_{x_0} centered at x_0 . Then it is singular with respect to \mathcal{L} . Indeed fix $A = \mathbb{R} \setminus \{x_0\}$, $B = \{x_0\}$, and observe that $\mathcal{L}(B) = 0$ and $\delta_{x_0}(A) = 0$.

Proposition 2.30. Let $f \geq 0$, measurable and such that $\int_{-M}^M f(x)dx < +\infty$ for all $M > 0$. Define the function

$$\nu_f : \mathcal{M} \rightarrow [0, +\infty] \quad \text{as} \quad \mu_f(A) = \int_A f(x)dx.$$

Then ν_f is a measure on $(\mathbb{R}^n, \mathcal{M})$, which is σ -finite and which is absolutely continuous with respect to \mathcal{L} . If $f \in L^1(\mathbb{R}^n)$ the measure is finite.

Proof. First of all we show that it is a measure. Observe that $f(x)\chi_\emptyset(x) = 0$ almost everywhere, then $\nu_f(\emptyset) = 0$. Let $A_i \in \mathcal{M}$ which are pairwise disjoint. Define the simple function $\phi_k(x) = \sum_{i=1}^k \chi_{A_i}(x)$. Note that $\lim_k \phi_k(x) = \chi_{\cup_i A_i}(x)$. Moreover $0 \leq f(x)\phi_k(x) \leq f(x)\phi_{k+1}(x)$ and so by the monotone convergence theorem we get

$$\lim_k \int_{\mathbb{R}^n} \phi_k(x)f(x)dx = \int_{\mathbb{R}^n} \lim_k \phi_k(x)f(x)dx.$$

Observe that

$$\begin{aligned} \lim_k \int_{\mathbb{R}^n} \phi_k(x)f(x)dx &= \lim_k \int_{\mathbb{R}^n} \sum_{i=1}^k \phi_i(x)f(x)dx = \lim_k \sum_{i=1}^k \int_{\mathbb{R}^n} \phi_i(x)f(x)dx \\ &= \lim_k \sum_{i=1}^k \int_{A_i} f(x)dx = \lim_k \sum_{i=1}^k \nu_f(A_i) = \sum_{i=1}^{+\infty} \nu_f(A_i) \end{aligned}$$

and

$$\int_{\mathbb{R}^n} \lim_k \phi_k(x)f(x)dx = \int_{\mathbb{R}^n} \chi_{\cup_i A_i}(x)f(x)dx = \nu_f(\cup_i A_i).$$

Therefore we get that ν_f is a measure.

Since $\nu_f(B(0, k)) = \int_{B(0, k)} f(x)dx < +\infty$ by assumption, then ν_f is σ -finite.

Finally, note that if $A \in \mathcal{M}$ and $\mathcal{L}(A) = 0$, this implies that $\chi_A(x) = 0$ almost everywhere. Therefore also $f(x)\chi_A(x) = 0$ almost everywhere, which implies $\nu_f(A) = 0$. □

Example 2.31. Let $f(x) = e^{-|x|^2}$. Then $f \in L^1(\mathbb{R}^n)$ and the measure ν_f is called the Gaussian measure. Note that it is a finite measure, and $\int_{\mathbb{R}^n} e^{-|x|^2} dx = \pi^{n/2}$, see [2, Prop. 2.53].

Theorem 2.32 (Lebesgue-Radon-Nikodym theorem). Let μ a measure on $(\mathbb{R}^n, \mathcal{M})$ which is σ -finite. Then there exist a unique $\nu \ll \mathcal{L}$ (absolutely continuous part) and a unique $\rho \perp \mathcal{L}$ (singular part) such that $\mu = \nu + \rho$.

Moreover there exists $f \geq 0$, measurable and such that $\int_A f(x)dx < +\infty$ for all $A \in \mathcal{M}$ with $\mathcal{L}(A) < +\infty$, for which $\nu = \nu_f$. f is called the **density** of ν , or the *Radon-Nikodym derivative* of ν and can be obtained (if the measure ν is regular) as $f(x) = \lim_{r \rightarrow 0} \frac{\nu(B(x, r))}{\mathcal{L}(B(x, r))}$.

Proof. For the proof we refer to [2, Section 3.2]. □

2.7 Distribution of a random variable

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $X : \Omega \rightarrow \mathbb{R}$ be a random variable (see Section 2.4). Then **the distribution** \mathbb{P}_X of X is the Borel measure induced on \mathbb{R} by X , defined as follows: for every $A \in \mathcal{B}(\mathbb{R})$,

$$\mathbb{P}_X(A) = \mathbb{P}(\{\omega \mid X(\omega) \in A\}).$$

\mathbb{P}_X is an absolutely continuous measure, iff X is a continuous random variable, and \mathbb{P}_X is a singular measure with respect to the Lebesgue measure, iff X is a discrete random variable. The distribution identifies the random variable, and often the random variables are described just in terms of their distributions. Some examples of widely used random variables/distributions:

- the **gamma distribution** with parameters a, b is an absolutely continuous measure with density $f(x) = \Gamma(a)^{-1} b^a x^{a-1} e^{-bx} \chi_{(0, +\infty)}(x)$
- the **chi-square distribution** is a gamma distribution with parameters $n/2, 1/2$,
- the **normal or Gaussian distribution** with parameters μ, σ is an absolutely continuous random variable, with density $f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$,
- the **standard normal distribution** is a normal distribution with parameters $0, 1$, that is an absolutely continuous measure with density $f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$,
- the **binomial distribution** of parameters n, p is a singular measure, and it is given by $\sum_{k=0}^n \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} \delta_k$ where δ_k is the Dirac measure centered at k ,
- the **Poisson distribution** of parameter λ is a singular measure, and it is given by $e^{-\lambda} \sum_{k=0}^{+\infty} \frac{\lambda^k}{k!} \delta_k$ where δ_k is the Dirac measure centered at k .

Definition 2.33. *The n -th-moment of a random variable X is given by $\mathbb{E}(X^n)$, more precisely*

- *if X is a continuous random variable (whose associated distribution has density f) then*

$$\mathbb{E}(X^n) = \int_{\mathbb{R}} x^n f(x) dx.$$

- *if X is a discrete random variable (taking values on \mathbb{Z}),*

$$\mathbb{E}(X^n) = \sum_{k \in \mathbb{Z}} k^n P(\omega \mid X(\omega) = k).$$

Note that $\mathbb{E}(X^n) < +\infty$ if and only if $\mathbb{E}(|X|^n) < +\infty$.

*We recall that the moment for $n = 1$, that is $\mathbb{E}(X)$, is called the **mean**, whereas $\mathbb{E}(X^2) - (\mathbb{E}(X))^2$ is called the **variance**.*

2.8 Problems

- Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a monotone function. Show that f is Lebesgue measurable.
- Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a Lebesgue measurable function and let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $f(x) = g(x)$ for almost every $x \in \mathbb{R}^n$. Show that g is Lebesgue measurable.
- Consider the right continuous increasing function on \mathbb{R}

$$F(x) = \begin{cases} x & x < 0 \\ x + 1 & x \geq 0. \end{cases}$$

Which is the Borel measure associated to this function?

- Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a positive function. Let $F(t)$ the repartition function of f . Recall that $\int_{\mathbb{R}^n} f(x) dx = \int_0^{+\infty} F(t) dt$. Show that for all $p > 1$

$$\int_{\mathbb{R}^n} |f(x)|^p dx = p \int_0^{+\infty} t^{p-1} F(t) dt.$$

So $f \in L^p(\mathbb{R}^n)$ if and only if $p \int_0^{+\infty} t^{p-1} F(t) dt < +\infty$.

(v) Let

$$f(x) = \begin{cases} \frac{1}{|x|} & |x| \leq 1, x \neq 0 \\ 0 & |x| > 1, x = 0. \end{cases}$$

For which p , is the function $f \in L^p(\mathbb{R}^n)$?

(vi) Let

$$f(x) = \begin{cases} \log\left(\frac{1}{|x|}\right) & 0 < |x| < 1 \\ 0 & |x| = 0. \end{cases}$$

Show that $f \in L^1(B(0, 1))$.

Is it true that the function f satisfies $f \in L^p(B(0, 1))$ for every $p \geq 1$?

Is it true that $f \in L^\infty(B(0, 1))$?

(vii) Let

$$f(x) = \begin{cases} \frac{1}{|x|} & |x| > 1 \\ 1 & |x| \leq 1. \end{cases}$$

For which p , is the function f in the space $L^p(\mathbb{R}^n)$?

3 Banach spaces

3.1 Banach spaces

Let X be a vectorial space on \mathbb{R} (this means that it is closed by summation and by multiplication by scalars, that is if $x, y \in X$, $\lambda, \mu \in \mathbb{R}$, then $\lambda x + \mu y \in X$).

Definition 3.1. A norm $\|\cdot\| : X \rightarrow [0, +\infty)$ is a function such that

- $\|x\| \geq 0$ for all $x \in X$ and $\|x\| = 0$ iff $x = 0$ (positivity);
- $\|\lambda x\| = |\lambda| \|x\|$ for all $x \in X, \lambda \in \mathbb{R}$ (homogeneity);
- $\|x + y\| \leq \|x\| + \|y\|$ (triangular inequality).

$(X, \|\cdot\|)$ is a normed space.

Example 3.2. On \mathbb{R}^n we may define the euclidean norm $|x| = \sqrt{x_1^2 + \dots + x_n^2}$.

A norm induces on X a metric structure on X in the following way.

Definition 3.3 (Metric structure and notion of convergence). Let $(X, \|\cdot\|)$ be a normed space. We define a distance between elements in X as

$$d(x, y) = \|x - y\|.$$

Note that this is a good definition, since it is positive, zero only if $x = y$, and satisfies the triangular inequality, that is $d(x, z) \leq d(x, y) + d(y, z)$ for all x, y, z .

We define the balls associated to this distance as follows: we fix a center $x_0 \in X$ and a radius $r > 0$ and we set

$$B(x_0, r) = \{x \in X \mid \|x - x_0\| < r\}.$$

A set $A \subseteq X$ is **open** if for all $x \in A$ there exists $r > 0$ such that $B(x, r) \subseteq A$. A set C is **closed** if $X \setminus C$ is open.

Let $(x_n)_n$ a sequence of element in X and $x \in X$. Then

$$\lim_n x_n = x \quad \text{iff} \quad \lim_{n \rightarrow +\infty} \|x_n - x\| = 0.$$

Proposition 3.4. The following are equivalent:

- i) C is closed
- ii) for every sequence (x_n) of elements in C such that there exists $x \in X$ with $\lim_n x_n = x$, there holds that $x \in C$.

Proof. Assume that C is closed and ii) is false. Then there exists (x_n) of elements in C such that $\lim_n x_n = x \notin C$. This implies that there exists $r > 0$ such that $B(x, r) \subseteq X \setminus C$. Therefore $x_n \notin B(x, r)$ for all n , which means that $\|x_n - x\| \geq r$ for all n , in contradiction with the fact that $\lim_n x_n = x$.

Assume that ii) holds and assume that C is not closed. So there exists $x \notin C$ such that for all $r > 0$ there holds that $B(x, r) \cap C \neq \emptyset$. Let $x_n \in C$ such that $x_n \in B(x, \frac{1}{n}) \cap C$. So $\|x_n - x\| < \frac{1}{n}$ and then $\lim_n x_n = x$. But this would imply $x \in C$. \square

Definition 3.5 (Banach space).

A sequence $(x_n)_n$ in X is a *Cauchy sequence* if $\lim_{n,m} \|x_n - x_m\| = 0$.

A normed space is called a *Banach space* if all the Cauchy sequences have limit in X .

Remark 3.6. Note that if $(x_n)_n$ is a sequence which converge to $x \in X$, then it is also a Cauchy sequence, since by triangular inequality $\|x_n - x_m\| \leq \|x_n - x\| + \|x - x_m\|$ and then $0 \leq \lim_{n,m \rightarrow +\infty} \|x_n - x_m\| \leq \lim_{m,n \rightarrow +\infty} (\|x_n - x\| + \|x - x_m\|) = 0$.

The viceversa is not always true. Let's think e.g. of the case $X = \mathbb{Q}$ and the euclidean norm. Define (x_n) as follows: $x_0 = 1$, $x_1 = 1,01$, $x_2 = 1,01001$, $x_3 = 1,010010001$, $x_4 = 1,01001000100001$ and so on, that is $x_n = 1,1010010001 \dots 1 \overset{n}{0} \dots 0 1$. It is easy to check that $x_n \in \mathbb{Q}$ for all n , that $x_n \rightarrow x$ (so $(x_n)_n$ is a Cauchy sequence, but this can also be checked directly) and that $x \notin \mathbb{Q}$. This implies that $(\mathbb{Q}, |\cdot|)$ is not a Banach space.

An important theorem in Banach spaces (more generally in complete metric spaces) is the **contraction lemma**, or **Banach-Caccioppoli theorem**:

Theorem 3.7. Let $(X, \|\cdot\|)$ a Banach space and $F : X \rightarrow X$ such that there exists $0 < a < 1$ for which

$$\|F(x) - F(y)\| \leq a\|x - y\| \quad \forall x, y \in X.$$

(F is a contraction) Then the map F admits a unique fixed point, that is a point such that $\bar{x} = F(\bar{x})$.

Proof. See problem 1 at the end of the chapter. □

3.2 Bounded linear operators

Definition 3.8. Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be two Banach space.

A **linear operator** is a map $T : X \rightarrow Y$ such that $T(\alpha x + \beta y) = \alpha T(x) + \beta T(y)$ for all $\alpha, \beta \in \mathbb{R}$, $x, y \in X$.

A **bounded operator** is a map $T : X \rightarrow Y$ such that

$$\|T\| = \sup_{\{x \in X \mid \|x\| \leq 1\}} \|Tx\| < +\infty.$$

If this quantity is finite, it is called the norm of T .

A **continuous operator** is a map $T : X \rightarrow Y$ such that

$$\lim_{n \rightarrow \infty} \|Tx_n - Tx\| = 0 \quad \text{for all sequences } x_n \text{ such that } \lim_{n \rightarrow \infty} x_n = x.$$

Proposition 3.9. A linear operator $T : X \rightarrow Y$ is continuous if and only if it is bounded.

Proof. Assume that T is bounded, then

$$\|Tx_n - Tx\| = \|T(x_n - x)\| = \|x_n - x\| \|T\| \leq \|x_n - x\| \|T\|.$$

Therefore if $\|x_n - x\| \rightarrow 0$, then also $\|Tx_n - Tx\| \rightarrow 0$.

Assume that T is continuous, and we want to prove that T is bounded. Assume by contradiction that it is not true. So for every $n \in \mathbb{N}$ there exists $x_n \in X$ such that $\|x_n\| = 1$ and $\|Tx_n\| \geq n$. Define $y_n = \frac{x_n}{n}$. Then $\|y_n\| = \frac{\|x_n\|}{n} = \frac{1}{n} \rightarrow 0$. This implies that $y_n \rightarrow 0$. Observe that by linearity $Ty_n = \frac{1}{n}Tx_n$ and then $\|Ty_n\| = \frac{1}{n}\|Tx_n\| \geq \frac{n}{n} = 1$. Therefore $y_n \rightarrow 0$ but $\|Ty_n\| \not\rightarrow 0$, in contradiction with continuity. □

Theorem 3.10. The set of all bounded linear operators between two Banach spaces X, Y , is a Banach space $\mathcal{B}(X, Y)$, with norm $\|T\|$ as defined above.

Proof. See [1, Theorem 2.12]. □

Example 3.11. Let $X = \mathbb{R}^n$ and $Y = \mathbb{R}^m$ both with the euclidean norm. Let $\mathbf{A} \in M_{m \times n}(\mathbb{R})$ be a $n \times m$ matrix. Then

$$Tx = \mathbf{A}x = \left(\sum_{j=1}^n a_{ij}x_j \right)_{i=1, \dots, m}$$

is a bounded linear operator from \mathbb{R}^n to \mathbb{R}^m .

Theorem 3.12 (Uniform boundedness principle, or Banach-Steinhaus theorem). *Let T_n be a sequence of bounded linear operators between the Banach spaces X and Y , that is $T_n \in \mathcal{B}(X, Y)$ for all n . Assume that for all $x \in X$ there exists $C_x \in \mathbb{R}$ such that $\sup_n \|T_n x\| \leq C_x$.*

Then there exists $C \in \mathbb{R}$ such that $\|T_n\| \leq C$ for all n .

In particular this implies that if the sequence $T_n x$ is convergent for every $x \in X$, then $Tx := \lim_n T_n x$ defines a bounded linear operator.

Proof. See [1, Theorem 4.1]. □

3.3 L^p spaces

We consider the L^p spaces defined in Section 2, Definition 2.27.

Definition 3.13. *Let $p > 1$. Then the conjugate exponent of p is the number $q > 1$ such that $1/p + 1/q = 1$. In particular the conjugate exponent of 2 is 2.*

We say that the conjugate exponent of 1 is $+\infty$.

Lemma 3.14 (Young inequality). *Let p, q be conjugate exponents. Then $xy \leq x^p/p + y^q/q$ for all $x, y \geq 0$.*

Proof. Fix $x > 0$ and consider $\sup_{y \geq 0} (xy - y^q/q)$. First of all observe that the supremum is actually a maximum, since $\lim_{y \rightarrow +\infty} xy - y^q/q = -\infty$. Differentiating in y , we get that the unique point where the derivative is 0 is given by $y = x^{1/(q-1)}$. This is the maximum. Therefore for all $y \geq 0$, $xy - y^q/q \leq x^{1+1/(q-1)} - x^{q/(q-1)}/q = x^p/p$, since $p = q/(q-1)$. □

Theorem 3.15 (Holder inequality). *Let $O \subseteq \mathbb{R}^n$ be an open set (it can also be $O = \mathbb{R}^n$), $p \in [1, +\infty]$ and q its conjugate exponent. Assume that $f \in L^p(O), g \in L^q(O)$. Then $f(x)g(x) \in L^1(O)$ and*

$$\int_O |f(x)g(x)| dx \leq \left(\int_O |f(x)|^p dx \right)^{1/p} \left(\int_O |g(x)|^q dx \right)^{1/q}.$$

Proof. Let $\tilde{f}(y) = |f(y)| \left(\int_O |f(x)|^p dx \right)^{-1/p}$ and $\tilde{g}(y) = |g(y)| \left(\int_O |g(x)|^q dx \right)^{-1/q}$. We apply the Young inequality to $\tilde{f}(y)$ and $\tilde{g}(y)$ and we get

$$|f(y)g(y)| \left(\int_O |f(x)|^p dx \right)^{-1/p} \left(\int_O |g(x)|^q dx \right)^{-1/q} \leq \frac{1}{p} \frac{|f(y)|^p}{\int_O |f(x)|^p dx} + \frac{1}{q} \frac{|g(y)|^q}{\int_O |g(x)|^q dx}$$

Integrating in O both sides we conclude

$$\frac{\int_O |f(x)g(x)| dx}{\left(\int_O |f(x)|^p dx \right)^{1/p} \left(\int_O |g(x)|^q dx \right)^{1/q}} \leq \frac{1}{p} + \frac{1}{q} = 1.$$

□

Corollary 3.16 (Minkowski inequality). *Let $f, g \in L^p(O)$, then*

$$\left(\int_O |f(x) + g(x)|^p dx \right)^{\frac{1}{p}} \leq \left(\int_O |f(x)|^p dx \right)^{\frac{1}{p}} + \left(\int_O |g(x)|^p dx \right)^{\frac{1}{p}}.$$

Proof. For $p = 1, \infty$, the inequality is straightforward. We consider the case $p \in (1, +\infty)$. First of all, we observe that if $f, g \in L^p$ the $f + g \in L^p$. This is due to the fact that

$$\frac{|f(x) + g(x)|^p}{2^p} = \left| \frac{f(x)}{2} + \frac{g(x)}{2} \right|^p \leq \frac{|f(x)|^p}{2} + \frac{|g(x)|^p}{2}$$

by the convexity of the function $r \mapsto r^p$ on $[0, +\infty)$ when $p \geq 1$. Now we observe that

$$|f(x) + g(x)|^p = |f(x) + g(x)||f(x) + g(x)|^{p-1} \leq |f(x)||f(x) + g(x)|^{p-1} + |g(x)||f(x) + g(x)|^{p-1}$$

and that $|f(x) + g(x)|^{p-1} \in L^q$ where $q = \frac{p}{p-1}$ is the conjugate exponent of p . Moreover

$$\int_O (|f(x) + g(x)|^{p-1})^q dx = \int_O |f(x) + g(x)|^p dx. \quad (3.1)$$

So by Holder inequality applied to f and $|f + g|^{p-1}$ we get

$$\int_O |f(x)| |f(x) + g(x)|^{p-1} dx \leq \left(\int_O |f(x)|^p dx \right)^{\frac{1}{p}} \left(\int_O |f(x) + g(x)|^p dx \right)^{\frac{p-1}{p}}$$

and analogously by Holder inequality applied to f and $|f + g|^{p-1}$ we get

$$\int_O |g(x)| |f(x) + g(x)|^{p-1} dx \leq \left(\int_O |g(x)|^p dx \right)^{\frac{1}{p}} \left(\int_O |f(x) + g(x)|^p dx \right)^{\frac{p-1}{p}}.$$

Integrating (3.1) and using the previous inequalities we get

$$\begin{aligned} \int_O |f(x) + g(x)|^p dx &\leq \left(\int_O |f(x)|^p dx \right)^{\frac{1}{p}} \left(\int_O |f(x) + g(x)|^p dx \right)^{\frac{p-1}{p}} \\ &\quad + \left(\int_O |g(x)|^p dx \right)^{\frac{1}{p}} \left(\int_O |f(x) + g(x)|^p dx \right)^{\frac{p-1}{p}} \\ &= \left(\int_O |f(x) + g(x)|^p dx \right)^{\frac{p-1}{p}} \left[\left(\int_O |f(x)|^p dx \right)^{\frac{1}{p}} + \left(\int_O |g(x)|^p dx \right)^{\frac{1}{p}} \right] \end{aligned}$$

from which we deduce the statement by dividing both sides by $\left(\int_O |f(x) + g(x)|^p dx \right)^{\frac{p-1}{p}}$. \square

Theorem 3.17. *Let $p \geq 1$.*

The spaces $L^p(O)$ are Banach spaces, with norm given by $\|f\|_p = \left(\int_O |f(x)|^p dx \right)^{1/p}$. The space $L^\infty(O)$ is a Banach space with norm given by $\|f\|_\infty = \inf\{c > 0 \mid \mathcal{L}\{x \mid |f(x)| \geq c\} = \emptyset\}$.

Proof. Proving that L^p is a vectorial space is an easy task only if $p = 1, +\infty$, otherwise it is a consequence of Minkowski inequality (which also gives that $\|\cdot\|_p$ satisfies the triangular inequality), which is a consequence of the Holder inequality. For the proof see [2, Section 6.1]. \square

A direct consequence of the Holder inequality is the following interpolation inequality.

Corollary 3.18 (Interpolation inequality). *Let $O \subseteq \mathbb{R}^n$ be an open set (it can also be $O = \mathbb{R}^n$), $p, r \in [1, +\infty]$ such that $p < r$. Assume that $f \in L^p(O) \cap L^r(O)$. Then $f \in L^s(O)$ for every $s \in [p, r]$ and moreover*

$$\|f\|_s = \|f\|_p^\alpha \|f\|_r^{1-\alpha} \quad \text{where } \alpha \in [0, 1] \text{ is such that } \frac{1}{s} = \frac{\alpha}{p} + \frac{1-\alpha}{r}.$$

Proof. Take $s \in (p, r)$ and $\alpha \in (0, 1)$ such that $\frac{1}{s} = \frac{\alpha}{p} + \frac{1-\alpha}{r}$. Since $1 = \frac{\alpha s}{p} + \frac{(1-\alpha)s}{r}$ we deduce that $\frac{p}{\alpha s} > 1$ and $\frac{r}{(1-\alpha)s} > 1$ are conjugate exponents.

Since $f \in L^p$, we get that $|f|^{\alpha s} \in L^{\frac{p}{\alpha s}}$ and moreover since $f \in L^r$ then $f^{(1-\alpha)s} \in L^{\frac{r}{(1-\alpha)s}}$. Therefore by the Holder inequality we get that $|f|^{\alpha s} |f|^{(1-\alpha)s} = |f|^s \in L^1$, which implies that $f \in L^s(O)$ and moreover

$$\|f\|_s^s = \int_O |f|^s dx \leq \left(\int_{\mathbb{R}} (|f|^{\alpha s})^{\frac{p}{\alpha s}} dx \right)^{\frac{\alpha s}{p}} \left(\int_{\mathbb{R}} (|f|^{(1-\alpha)s})^{\frac{r}{(1-\alpha)s}} dx \right)^{\frac{(1-\alpha)s}{r}} = \|f\|_p^{\alpha s} \|f\|_r^{(1-\alpha)s}.$$

\square

Another consequence of the Holder inequality is the following:

Corollary 3.19. *Let O be an open set with $\mathcal{L}(O) < +\infty$. Then $L^p(O) \subseteq L^r(O)$ for every $1 \leq r \leq p$, and moreover $\|f\|_r \leq \|f\|_p \mathcal{L}(O)^{\frac{p-r}{pr}}$.*

Proof. Fix $p > 1$ and $f \in L^p(O)$. We want to prove that $f \in L^1(O)$. Note that since $\mathcal{L}(O) < +\infty$, then $\chi_O \in L^q(\mathbb{R}^n)$ for every q , so in particular it is in $L^q(O)$ for q conjugate exponent of p . By Holder inequality we get

$$\|f\|_1 \leq \|f\|_p \mathcal{L}(O)^{1/q} = \|f\|_p \mathcal{L}(O)^{\frac{p-1}{p}}$$

which give the conclusion of the theorem for $r = 1$. The case $r \in (1, p)$ is obtained just using the interpolation inequality, proved in the previous corollary: indeed $\frac{1}{r} = \frac{\alpha}{p} + 1 - \alpha$, with $\alpha = \frac{p(r-1)}{r(p-1)}$ and then

$$\|f\|_r \leq \|f\|_p^\alpha \|f\|_1^{1-\alpha} \leq \|f\|_p^\alpha \|f\|_p^{1-\alpha} \mathcal{L}(O)^{\frac{p-1}{p}(1-\alpha)} = \|f\|_p \mathcal{L}(O)^{\frac{p-r}{pr}}.$$

□

Finally we present an important example of linear bounded operators from L^p to \mathbb{R} .

Example 3.20. Let $g \in L^q(\mathbb{R}^n)$ with $q \geq 1$. Consider the following operator $T : L^p(\mathbb{R}^n) \rightarrow \mathbb{R}$, where p is the conjugate exponent of q , defined as

$$Tf = \int_{\mathbb{R}^n} f(x)g(x)dx.$$

It is immediate to check that it is linear. Moreover, by Holder inequality we get, for all $f \in L^p(\mathbb{R}^n)$ with $\|f\|_p \leq 1$,

$$|Tf| = \left| \int_{\mathbb{R}^n} f(x)g(x)dx \right| \leq \int_{\mathbb{R}^n} |f(x)g(x)|dx \leq \|f\|_p \|g\|_q \leq \|g\|_q.$$

Therefore T is a bounded operator, with norm $\|T\| \leq \|g\|_q$.

Define now $f_g(x) = |g(x)|^{q/p} \|g\|_q^{-q/p} \frac{g(x)}{|g(x)|}$. Then $f_g \in L^p(\mathbb{R}^n)$ and $\|f_g\|_p = \|g\|_q^{q/p} \|g\|_q^{-q/p} = 1$. We compute, recalling that $q/p + 1 = q$

$$Tf_g = \|g\|_q^{q/p} \int_{\mathbb{R}^n} |g(x)|^{q/p+1} dx = \|g\|_q^{q/p} \int_{\mathbb{R}^n} |g(x)|^q dx = \|g\|_q^{q-q/p} = \|g\|_q.$$

Therefore $\|T\| = \|g\|_q$.

3.4 Spaces of random variables with finite moments

We fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and we consider the random variables $X : \Omega \rightarrow \mathbb{R}$. We introduce the spaces of random variables with finite p -moment (see definition in Section 2.7)

$$M^p = \{X \text{ random variable } \mathbb{E}(|X|^p) < +\infty\}$$

and we $\|X\|_p = (\mathbb{E}(|X|^p))^{1/p}$.

First of all we have the following Holder inequality and Minkowski inequality

Proposition 3.21. *Let $X \in M^p$ and $Y \in M^q$, with q conjugate exponent of p , then*

$$\mathbb{E}(|XY|) \leq \mathbb{E}(|X|^p)^{1/p} \mathbb{E}(|Y|^q)^{1/q}.$$

Moreover if $X, Z \in M^p$, then

$$\mathbb{E}(|X + Z|^p)^{1/p} \leq \mathbb{E}(|X|^p)^{1/p} + \mathbb{E}(|Z|^p)^{1/p}$$

Proof. It is sufficient to apply the Young inequality to $|X|\mathbb{E}(|X|^p)^{-1/p}$ and to $|Y|\mathbb{E}(|Y|^q)^{-1/q}$ and proceed as for L^p spaces (instead of integrating in \mathcal{O} one needs to take the average of both side of the inequality). The proof of Minkowski also follows again as for the case of L^p spaces. \square

Theorem 3.22. *The space M^p with the norm $\|X\|_p$ for $p \in [1, +\infty)$ is a Banach space.*

We say that $X_n \rightarrow X$ in M^1 if $\mathbb{E}(|X_n - X|) \rightarrow 0$, that is **convergence in mean** and $X_n \rightarrow X$ in M^2 if $\mathbb{E}((X_n - X)^2) \rightarrow 0$, that is **convergence in the mean square convergence**.

Similarly as for $L^p(\mathcal{O})$ spaces, where $\mathcal{L}(\mathcal{O}) < +\infty$ (see Corollary 3.19), the spaces M^p are decreasing as we will show. First of all we recall the Jensen inequality:

Lemma 3.23 (Jensen's inequality). *Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a convex function, then for every random variable X*

$$\mathbb{E}(g(X)) \geq g(\mathbb{E}(X)).$$

Theorem 3.24. *There holds that $M^k \subseteq M^n$ for every $1 \leq n \leq k$. Moreover if $X \in M^k$ then $(\mathbb{E}(|X|^n))^{\frac{1}{n}} \leq (\mathbb{E}(|X|^k))^{\frac{1}{k}}$ for all $n \leq k$.*

Proof. Let $1 \leq n \leq k$, $g(x) = |x|^{\frac{k}{n}}$. Since $\frac{k}{n} \geq 1$, the function g is convex. Let $X \in M^k$ and we apply Jensen's inequality to the random variable $|X|^n$, observing that $g(|X|^n) = |X|^k$,

$$\mathbb{E}(|X|^k) = \mathbb{E}(g(|X|^n)) \geq g(\mathbb{E}(|X|^n)) = (\mathbb{E}(|X|^n))^{\frac{k}{n}}.$$

\square

3.5 Modes of convergence and compactness theorems

Note that $f_n \rightarrow f$ in $L^p(\mathcal{O})$ means that $\lim_n \int_{\mathcal{O}} |f_n(x) - f(x)|^p dx = 0$. Moreover $f_n \rightarrow f$ in L^∞ if $\lim_n \sup_{\mathcal{O}} |f_n - f| = 0$. **The convergence in L^1 is also called convergence in mean, and the convergence in L^2 is called convergence in mean square.**

Definition 3.25. $f_n \rightarrow f$ almost everywhere if $\lim_n f_n(x) = f(x)$ for almost every x .

It is not always true that convergence almost everywhere is sufficient for convergence in L^p as the following example shows.

Example 3.26. Let

$$f_n(x) = \begin{cases} 0 & x \geq \frac{1}{n} \\ n - nx & 0 \leq x \leq \frac{1}{n} \\ n + nx & -\frac{1}{n} \leq x \leq 0 \\ 0 & x \leq -\frac{1}{n}. \end{cases}$$

Then $\lim_n f_n(x) = 0$ for almost every x , but $\int_{\mathbb{R}} f_n(x) dx = 2 \neq 0$.

Theorem 3.27. *Let $p \geq 1$, $(f_n)_n, f \in L^p(\mathcal{O})$ and assume that $f_n \rightarrow f$ almost everywhere.*

If there exists $g \in L^p(\mathcal{O})$ such that $|f_n(x)| \leq g(x)$ for almost every x and every n then $\lim_n f_n = f$ in $L^p(\mathcal{O})$.

If $\lim_n f_n = f$ in $L^p(\mathcal{O})$, then up to passing to a subsequence $f_n \rightarrow f$ almost everywhere.

If $\lim_n f_n = f$ in $L^p(\mathcal{O})$, then $f_n \rightarrow f$ in measure.

Proof. The first part is the Lebesgue dominated convergence theorem, see [2, Theorem 2.24].

The second part is proven in [2, Corollary 2.32].

The third part is a consequence of the Chebycheff inequality (see Problem ii). Indeed

$$\forall \varepsilon > 0 \quad \mathcal{L}(\{x \in \mathbb{R}^n \mid |f_n(x) - f(x)| > \varepsilon\})^{\frac{1}{p}} \leq \frac{1}{\varepsilon} \|f_n - f\|_p.$$

\square

By the Holder inequality we can multiply functions in L^p by functions in L^q . This gives another notion of convergence.

Definition 3.28 (Weak convergence). *Let $1 < p < +\infty$. Given $(f_n)_n, f \in L^p(O)$, we say that $f_n \rightharpoonup f$ (weakly) in $L^p(O)$ if for all $g \in L^q(O)$, with q the conjugate exponent of p , ($q = +\infty$ if $p = 1$) there holds*

$$\lim_n \int_O f_n(x)g(x)dx = \int_O f(x)g(x)dx.$$

Given $(f_n)_n, f \in L^\infty(O)$, we say that $f_n \rightharpoonup^ f$ (weakly star) in $L^\infty(O)$ if for all function $g \in L^1(O)$, there holds*

$$\lim_n \int_O f_n(x)g(x)dx = \int_O f(x)g(x)dx.$$

Given $(f_n)_n, f \in L^1(O)$, we say that $f_n \rightharpoonup f$ (weakly) in $L^1(O)$ if for all continuous and bounded functions $g : O \rightarrow \mathbb{R}$, there holds

$$\lim_n \int_O f_n(x)g(x)dx = \int_O f(x)g(x)dx.$$

*If f_n are densities of continuous random variables X_n , this convergence is also called **convergence in distribution** of X_n .*

Proposition 3.29. *If f_n converge to f in L^p then it also converge weakly in L^p , whereas the viceversa is not true.*

Proof. The statement is a consequence of Holder inequality: let $f_n, f \in L^p$ and $g \in L^q$ (for $p > 1$) or g continuous and bounded (if $p = 1$), then

$$\left| \int_O (f_n(x) - f(x))g(x)dx \right| \leq \int_O |f_n(x) - f(x)||g(x)|dx \leq \|f_n - f\|_p \|g\|_q.$$

Therefore if $\|f_n - f\|_p \rightarrow 0$, then $\int_O (f_n(x) - f(x))g(x)dx \rightarrow 0$. □

The main examples of sequence of functions which are converging weakly but not strongly are rapidly oscillating functions.

Example 3.30. [Weak convergence of periodic functions] Let $f(x)$ be a continuous periodic function (e.g, $f(x) = \sin x$) in \mathbb{R} with period T .

Define $f_n(x) = f(nx)$ (note that this is a periodic function with period T/n , so as $n \rightarrow +\infty$ this is more and more oscillating).

Then for every $O \subseteq \mathbb{R}^n$ borelian bounded set

$$f_n \rightharpoonup \frac{1}{T} \int_0^T f(x)dx \quad \text{in } L^p(O) \text{ for all } 1 \leq p < +\infty$$

and moreover

$$f_n \rightharpoonup^* \frac{1}{T} \int_0^T f(x)dx \quad \text{in } L^\infty(O).$$

For the proof see [1, Example 5.16].

Intuitively weak convergence is convergence of mean values, in the following sense.

Theorem 3.31 (Characterization of weak convergence). *Let $f_n \in L^p(O)$ for $p \in (1, +\infty]$ such that there exists $C > 0$ such that $\|f_n\|_p \leq C$ for all n . They are equivalent*

- (i) $f_n \rightharpoonup f$ (weakly) in $L^p(O)$ (or weakly star if $p = +\infty$)

(ii) for every Borel set $A \subseteq O$ with $|A| > 0$ there holds

$$\frac{1}{|A|} \int_A f_n(x) dx \rightarrow \frac{1}{|A|} \int_A f(x) dx.$$

Let $f_n \in L^1(O)$ such that there exists $C > 0$ such that $\|f_n\|_1 \leq C$ for all n . Then they are equivalent

(i) $f_n \rightarrow f$ in $L^1(O)$

(ii) for all open sets $A \subseteq O$ with $|A| > 0$

$$\lim_n \frac{1}{|A|} \int_A f_n(x) dx \geq \frac{1}{|A|} \int_A f(x) dx$$

and for all compact sets $C \subseteq O$ with $|C| > 0$

$$\lim_n \frac{1}{|C|} \int_C f_n(x) dx \leq \frac{1}{|C|} \int_C f(x) dx.$$

We get the following important compactness result.

Theorem 3.32 (Compactness). *Let $f_n \in L^p(O)$ for $p \in (1, +\infty]$ such that there exists $C > 0$ such that $\|f_n\|_p \leq C$ for all n . Then up to passing to a subsequence $f_n \rightarrow f$ (weakly) in $L^p(O)$ (or weakly star if $p = +\infty$).*

Let $f_n \in L^1(O)$ such that there exists $C > 0$ such that $\|f_n\|_1 \leq C$ for all n and moreover for all sequences $E_k \subseteq O$ such that $\chi_{E_k} \rightarrow 0$ in $L^1(O)$, there holds $\lim_k \sup_n \int_{E_k} f_n(x) dx \rightarrow 0$ (f_n are equiintegrable). Then up to passing to a subsequence $f_n \rightarrow f$ (weakly) in $L^1(O)$.

Theorem 3.33 (Prokhorov's theorem). *Let f_n are densities of continuous random variables $X_n : \Omega \rightarrow \mathbb{R}^d$, which are **tight** in the following sense: for every $\varepsilon > 0$ there exists a compact set $K_\varepsilon \subseteq \mathbb{R}^d$ such that $\mathbb{P}\{\omega, X_n(\omega) \in K_\varepsilon\} \geq 1 - \varepsilon$ for all n . Then, there exists f density of a continuous random variable X such that, up to a subsequence, $f_n \rightarrow f$ weakly (or equivalently $X_n \rightarrow X$ in distribution).*

3.6 Problems

(i) Let $(X, \|\cdot\|)$ a Banach space and $F : X \rightarrow X$ such that there exists $0 < a < 1$ for which

$$\|F(x) - F(y)\| \leq a\|x - y\| \quad \forall x, y \in X.$$

(F is a contraction)

(a) Show that the map F is continuous.

(b) Let $x_0 \in X$. Define $x_1 = F(x_0)$, $x_2 = F(x_1)$ and so on $x_n = F(x_{n-1})$. Prove that

$$\|x_n - x_{n+1}\| \leq a^n \|x_0 - x_1\|.$$

Deduce that $(x_n)_n$ is a Cauchy sequence.

(c) Let $\bar{x} = \lim_n x_n$, where (x_n) has been defined in the previous step. Show that $F(\bar{x}) = \bar{x}$. So, \bar{x} is a fixed point of F .

(d) Show that the map F admits a unique fixed point, that is a point such that $\bar{x} = F(\bar{x})$.

This is called **Banach-Caccioppoli theorem**.

(ii) Let $f \in L^p(\mathbb{R}^n)$ and $\alpha > 0$. Prove that

$$\mathcal{L}(\{x \in \mathbb{R}^n \mid |f(x)| > \alpha\})^{\frac{1}{p}} \leq \frac{1}{\alpha} \|f\|_p.$$

This is called **Chebycheff inequality**.

(iii) Let $f \in L^\infty(O) \cap L^p(O)$ for some $p > 1$. Our aim is to show that $\lim_{q \rightarrow +\infty} \|f\|_q = \|f\|_\infty$.

(a) Show that $f \in L^q(O)$ for every $q \geq p$.

(b) Prove that $\lim_{q \rightarrow +\infty} \|f\|_q \leq \|f\|_\infty$.

(c) By using the Chebycheff inequality prove that for every $a > 0$ and every $q \geq p$

$$a \mathcal{L}(\{x \in \mathbb{R}^n \mid |f(x)| > a\})^{\frac{1}{q}} \leq \|f\|_q.$$

Observe that if $\mathcal{L}(\{x \in \mathbb{R}^n \mid |f(x)| > a\}) \neq 0$, then $a \leq \lim_{q \rightarrow +\infty} \|f\|_q$.

(d) Using the previous point show that $\lim_{q \rightarrow +\infty} \|f\|_q \geq \|f\|_\infty$ and therefore by point 2, $\lim_{q \rightarrow +\infty} \|f\|_q = \|f\|_\infty$.

(iv) Prove that if $f \in L^2(-1, 1)$ then $f \in L^1(-1, 1)$ and moreover

$$\|f\|_1 \leq \sqrt{2} \|f\|_2.$$

Provide an example of a function $f \in L^1(-1, 1)$ such that $f \notin L^2(-1, 1)$.

(v) Let $f_n \in L^p(O)$ with $p > 1$ such that $f_n \rightharpoonup f$ weakly in L^p . Show that there exists $C > 0$ such that $\|f_n\|_p \leq C$ for all n .

(vi) Consider the following operator $T : L^2(0, 2) \rightarrow L^2(0, 2)$ defined as

$$Tf(x) = \int_0^x f(y) dy.$$

Show that this is a bounded continuous operator.

Hint Recall the **Jensen inequality**:

$$\left(\frac{1}{b-a} \int_a^b f(x) dx \right)^2 \leq \frac{1}{b-a} \int_a^b f(x)^2 dx.$$

(vii) Let $O \subseteq \mathbb{R}^n$ be a open set, $f_k, f \in L^p(O)$, for all $k \in \mathbb{N}$.

(a) Show that if $f_k \rightarrow f$ (strongly) in $L^p(O)$ then $\lim_k \|f_k\|_p = \|f\|_p$.

(b) Show that if $f_k \rightharpoonup f$ (weakly) in $L^p(O)$ then $\liminf_k \|f_k\|_p \geq \|f\|_p$.

(viii) Let $O \subseteq \mathbb{R}^n$ be a open set, let $p \geq 1$ and $q = \frac{p}{p-1}$ its conjugate exponent.

Let $f_k, f \in L^p(O)$, for all $k \in \mathbb{N}$, $g_k, g \in L^q(O)$, for all $k \in \mathbb{N}$.

(a) Assume $f_k \rightarrow f$ (strongly) in $L^p(O)$ and $g_k \rightarrow g$ (strongly) in $L^q(O)$.

Show that $f_k g_k \rightarrow f g$ (strongly) in $L^1(O)$.

(b) Assume $f_k \rightarrow f$ (strongly) in $L^p(O)$ and $g_k \rightharpoonup g$ (weakly) in $L^q(O)$ (this implies in particular that $\|g_k\|_q \leq M$).

Show that $f_k g_k \rightarrow f g$ (strongly) in $L^1(O)$.

(ix) Let $O \subseteq \mathbb{R}^n$ be a open set, and $f_k, f \in L^2(O)$, for all $k \in \mathbb{N}$ such that $f_k \rightharpoonup f$ weakly in $L^2(O)$.

Show that if $\lim_k \|f_k\|_2 = \|f\|_2$, then $f_k \rightarrow f$ strongly in $L^2(O)$.

4 Hilbert spaces

4.1 Hilbert spaces

Hilbert spaces are spaces where it is possible to define the notions of length and orthogonality, which allow to work with the elements geometrically, as if they were vectors in Euclidean space. First of all we recall some basic definitions.

Definition 4.1. A set X is a vector space on \mathbb{R} (a real vector space) if it is a set equipped with two operations, vector addition (which allows to add two vectors $x, y \in X$ to obtain another vector $x + y \in X$) and scalar multiplication (which allows us to “scale” a vector $x \in X$ by a real number c to obtain a vector $cx \in X$). Moreover we require that X contains a neutral element for the vector addition, that is an element $0 \in X$ such that $0 + x = x$ for every $x \in X$ and $x - x = 0$.

A scalar product on X is a function $(\cdot, \cdot) : X \times X \rightarrow \mathbb{R}$ such that

- $(x, x) \geq 0$ for all x and $(x, x) = 0$ iff $x = 0$;
- it is symmetric $(x, y) = (y, x)$ for all $x, y \in X$;
- it is linear, that is $(\alpha x + \beta y, z) = \alpha(x, z) + \beta(y, z)$ for all $x, y, z \in X, \alpha, \beta \in \mathbb{R}$.

We associate to a scalar product a norm in this way $\|x\| = \sqrt{(x, x)}$.

Proposition 4.2. The function $\|\cdot\| : X \rightarrow [0, +\infty)$ defined as $\|x\| = \sqrt{(x, x)}$ is a norm. Moreover the scalar product is continuous, that is if $x_n \rightarrow x$ in X and $y \in X$, then $(x_n, y) \rightarrow (x, y)$ in \mathbb{R} .

Proof. Positivity and homogeneity are obvious. To prove the triangle inequality one first need to prove the Cauchy Schwartz inequality $|(x, y)| \leq \|x\|\|y\|$. See [1, Theorem 5.1].

The continuity is an easy consequence of the Cauchy Schwartz inequality:

$$|(x_n - x, y)| \leq \|x_n - x\|\|y\|.$$

□

Definition 4.3 (Hilbert space). A space X with a scalar product which induces on X a norm such that X is a Banach space is called Hilbert space.

Proposition 4.4 (Parallelogram identity). For every $x, y \in H$, there holds

$$\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2.$$

Proof. By definition and by linearity and symmetry of the scalar product $\|x + y\|^2 = (x + y, x + y) = (x, x) + 2(x, y) + (y, y) = \|x\|^2 + 2(x, y) + \|y\|^2$, and $\|x - y\|^2 = (x - y, x - y) = \|x\|^2 - 2(x, y) + \|y\|^2$. It is sufficient to sum. □

Example 4.5. In \mathbb{R}^n we define the scalar product $(x, y) = x_1y_1 + x_2y_2 + \dots + x_ny_n$. The euclidean norm is the norm associated to this scalar product. So \mathbb{R}^n with this scalar product is a Hilbert space. This is the basic example of Hilbert space of finite dimension.

4.2 Orthogonality and projections in Hilbert spaces

Definition 4.6 (Orthogonal space). We say that $x, y \in X$ are orthogonal if $(x, y) = 0$.

If $S \subseteq X$ is a subset of X , we define the orthogonal subspace

$$S^\perp = \{x \in X \mid (x, s) = 0 \forall s \in S\}.$$

This a vectorial subspace of X .

Theorem 4.7 (Orthogonal projection). *Let $V \subseteq H$ be a closed subspace of a Hilbert space, $V \neq \{0\}$ and let $h \in H$.*

Then there exists a unique element $v \in V$ at minimal distance from h , that is such that $\|h - v\| = \min_{w \in V} \|h - w\|$. Moreover there exists a unique element $s \in V^\perp$ such that $h = v + s$.

The map $Pr_V : H \rightarrow V$ which associate $h \rightarrow v$ is called the orthogonal projection of H in V and it is a bounded linear operator of norm 1.

Proof. We consider the minimization problem $\min_{w \in V} \|h - w\|$ and we show that it admits a solution which is unique. Since $\|h - w\| \geq 0$ we get that $\inf_{w \in V} \|h - w\| = \delta \geq 0$. Let $v_n \in V$ such that $\delta \leq \|v_n - h\| \leq \delta + 1/n$. Then $(v_n)_n$ is a Cauchy sequence, since by parallelogram identity and linearity

$$\|v_n - v_m\|^2 = 2\|v_n - h\|^2 + 2\|v_m - h\|^2 - \|(v_n + v_m) - 2h\|^2 \leq 2(\delta + 1/n)^2 + 2(\delta + 1/m)^2 - 4\|h - (v_n + v_m)/2\|^2.$$

We conclude by recalling that since $(v_n + v_m)/2 \in V$ then $\|h - (v_n + v_m)/2\| \geq \delta$,

$$\|v_n - v_m\|^2 \leq 2(\delta + 1/n)^2 + 2(\delta + 1/m)^2 - 4\delta^2 = 4\delta/n + 4\delta/m + 1/n^2 + 1/m^2 \rightarrow 0 \quad \text{as } n, m \rightarrow +\infty.$$

Since H is a Banach space there exists $v \in H$ such that $\lim_n v_n = v$ and since V is closed then $v \in V$. By continuity, we conclude that $\|v - h\| = \delta = \inf_{w \in V} \|h - w\|$. v is the unique minimizer. Indeed if it were not the case, there would exists $v' \in V$ with $\|v - h\| = \|v' - h\| = \delta$. By parallelogram identity

$$\|v - v'\|^2 = 2\|v - h\|^2 + 2\|v' - h\|^2 - 4\|(v + v')/2 - h\|^2 \leq 2\delta^2 + 2\delta^2 - 4\delta^2 = 0$$

which implies $\|v - v'\| = 0$.

Let $w \in V$. We claim that $(h - v, w) = 0$. Since v is the point at minimum distance, then the function $\lambda \rightarrow \|h - v + \lambda w\|^2$ has minimum in $\lambda = 0$. Differentiating the function in λ it should be that the derivative in 0 is 0. $\frac{\|h - v + \lambda w\|^2}{d\lambda} = \frac{(h - v + \lambda w, h - v + \lambda w)}{d\lambda} = 2(h - v, w)$. Therefore $(h - v, w) = 0$. This means that $h - v \in V^\perp$.

Let $v = Pr_V(h)$, $v' = Pr_V(h')$ and let $\alpha, \beta \in \mathbb{R}$. Then $\alpha v + \beta v' \in V$ and $\alpha v + \beta v' - \alpha h - \beta h' \in V^\perp$. Therefore by uniqueness $Pr_V(\alpha h + \beta h') = \alpha v + \beta v'$. Then Pr_V is linear. Moreover since $(Pr_V h - h, Pr_V h) = 0$,

$$\|h\|^2 = \|h - Pr_V h + Pr_V h\|^2 = (h - Pr_V h + Pr_V h, h - Pr_V h + Pr_V h) = \|h - Pr_V h\|^2 + \|Pr_V h\|^2.$$

This implies that for all h with $\|h\| \leq 1$, $\|Pr_V h\|^2 = \|h\|^2 - \|h - Pr_V h\|^2 \leq 1$. So Pr_V is bounded. Moreover if $h \in V$, then $Pr_V h = h$. Therefore $\|Pr_V\| = 1$. □

Definition 4.8 (Orthonormal set). *A set $\{u_i, i \in I\}$ of elements in H is an orthonormal set if $\|u_i\| = 1$ for all i and $(u_i, u_j) = 0$ for all $i \neq j$.*

Proposition 4.9. *Let $\{u_i, i \in I\}$ be a orthonormal set. Then the following are equivalent*

- if $(x, u_i) = 0$ for all i , then $x = 0$
- $\|x\|^2 = \sum_i |(x, u_i)|^2$ for all $x \in H$,
- for all $x \in H$, $x = \sum_i (x, u_i) u_i$, (where the convergence is with respect to the norm of H).

An orthonormal set for which one of the previous conditions hold is called an orthonormal basis. Every Hilbert space admits a orthonormal basis.

Proof. See [2, Proposition 5.28]. □

Definition 4.10 (Separable space). *H is separable if it admits a countable orthonormal basis.*

Theorem 4.11 (Computation of the orthogonal projection). *Let V be a closed subspace of H and let $\{v_i, i \in I\}$ be an orthonormal basis of V . Then for all $h \in H$,*

$$Pr_V(h) = \sum_{i \in I} (h, v_i) v_i.$$

Proof. See [1, Theorem 5.10]. □

4.3 Hilbert space of random variables

We fix a probability space $(\Omega, \mathbb{P}, \mathcal{F})$ and we define the space

$$M^2 = \{X : \Omega \rightarrow \mathbb{R} \mid X \text{ random variable with } \mathbb{E}(X^2) < +\infty\}.$$

Recall that X is a random variable if $X^{-1}(A) \in \mathcal{F}$ for every $A \in \mathcal{B}$ (so for every A in the σ -algebra of Borel sets. Given X random variable, we define $\sigma(X) \subseteq \mathcal{F}$, that is **the σ -algebra generated by X** , as the minimal σ - algebra contained in \mathcal{F} which contains all the elements $X^{-1}(A) = \{\omega \in \Omega \mid X(\omega) \in A\}$ for every $A \in \mathcal{B}$. So it is the minimal σ -algebra which assures that X is measurable.

Note that if X is a constant random variable, so $X(\omega) = c$ for all $\omega \in \Omega$, then $X^{-1}(A) = \Omega$ if $c \in A$, and $X^{-1}(A) = \emptyset$ if $c \notin A$. So in this case $\sigma(X) = \{\emptyset, \Omega\}$, which is the minimal possible σ -algebra.

We define on M^2 the scalar product

$$(X, Y) = \mathbb{E}(XY)$$

and the induced norm is

$$\|X\| = \sqrt{\mathbb{E}(X^2)}.$$

It is possible to prove that M^2 with this norm and this scalar product is a Hilbert space. Observe that, as we did for L^p spaces, we are actually considering class of equivalence of random variables, since we are identifying two random variables X, Y such that $\mathbb{P}(\omega \mid X(\omega) = Y(\omega)) = 1$.

Given a subspace $S \subseteq M^2$ we consider the σ -algebra \mathcal{S} generated by S as the minimal σ -algebra which contains $\sigma(X)$ for all the random variables $X \in S$. We define the **conditional expectation of X given \mathcal{S}** as the orthogonal projection of X in S as defined and characterized in Theorem 4.7, that is

$$\mathbb{E}(X|\mathcal{S}) = Pr_S(X),$$

or equivalently

$$\mathbb{E}(X - \mathbb{E}(X|\mathcal{S}))^2 = \min_{Z \in S} \mathbb{E}(X - Z)^2$$

Remark 4.12 (Linear least square estimation). Given $X \in M^2$ we want to find for all $Y, a, b \in \mathbb{R}$ for which it is minimal $\mathbb{E}((Y - a - bX)^2)$. So, the **linear least square estimator** is given by

$$L(Y|X) = \mathbb{E}(X|\sigma(X)) = a + bX,$$

where a, b are the optimal values which minimize $\mathbb{E}((Y - a - bX)^2)$. This problem can be restated exactly as a projection problem: we define S as the space generated by $X, 1$ in M^2 , that is $S = \{Z = aX + b \in M^2, a \in \mathbb{R}, b \in \mathbb{R}\}$ and we want to find $Pr_S(Y)$.

In order to solve the problem, first of all we choose an orthonormal basis of S . A basis of S is given by $\{1, X\}$. Observe that if $\mathbb{E}(X) = (X, 1) \neq 0$, we have that X and 1 are not orthogonal, so we substitute X with the element $X - \mathbb{E}(X)$ which is orthogonal to 1 . Moreover we have to normalize this element by choosing $c \in \mathbb{R}$ such that $c^2 \mathbb{E}(X - \mathbb{E}(X))^2 = 1$. Since $\mathbb{E}(X - \mathbb{E}(X))^2 = \mathbb{E}(X^2) - (\mathbb{E}(X))^2 = Var(X)$, it is sufficient to choose $c = \sqrt{Var(X)}$. Therefore an orthonormal basis of S is given by $1, \frac{X - \mathbb{E}(X)}{\sqrt{Var(X)}}$. Recalling Theorem 4.11, we get

$$Pr_S(Y) = (Y, 1)1 + \left(Y, \frac{X - \mathbb{E}(X)}{\sqrt{Var(X)}} \right) \frac{X - \mathbb{E}(X)}{\sqrt{Var(X)}}.$$

So the linear least square estimator coincides with

$$L(Y|X) = \mathbb{E}(Y) + \frac{Cov(X, Y)}{Var(X)}(X - \mathbb{E}(X)).$$

Finally we compute the average error

$$\begin{aligned}\mathbb{E}(Y - L(Y|X))^2 &= \text{Var}(Y) + \frac{\text{Cov}^2(X, Y)}{\text{Var}^2(X)} \text{Var}X - 2 \frac{\text{Cov}(X, Y)}{\text{Var}(X)} \text{Cov}(X, Y) \\ &= \text{Var}Y - \frac{\text{Cov}^2(X, Y)}{\text{Var}(X)} = \frac{\text{Var}(Y)\text{Var}(X) - \text{Cov}^2(X, Y)}{\text{Var}(X)}.\end{aligned}$$

4.4 The Hilbert spaces L^2 and H^1

Let us fix an open set $O \subseteq \mathbb{R}^d$ and consider the Banach space $L^2(O)$ as defined in Definition 2.27. Note that by Holder inequality Theorem 3.15, if $f, g \in L^2(O)$, then $fg \in L^1(O)$ and then we may define a scalar product on $L^2(O)$ as follows:

$$(f, g) = \int_O f(x)g(x)dx.$$

It is immediate to check that actually it is a scalar product and moreover that $\|f\|_2 = (f, f)^{1/2}$. Therefore $L^2(O)$ is a Hilbert space.

Example 4.13 (Fourier serie). Let $L^2(-\pi, \pi)$. Then it is possible to show that an orthonormal basis of this space is given by $\{\frac{1}{\sqrt{2\pi}}, \frac{\cos(nx)}{\sqrt{\pi}}, \frac{\sin(nx)}{\sqrt{\pi}}, n \in \mathbb{N}\}$. By Parseval theorem every function $f \in L^2(-\pi, \pi)$ can be written as

$$f(x) = a_0 + \sum_{n=1}^{+\infty} a_n \cos nx + \sum_{n=1}^{+\infty} b_n \sin nx,$$

where

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)dx \quad a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx, \quad b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx.$$

This is called the Fourier serie of f .

The equality holds in the sense that $\lim_N \|f - a_0 - \sum_{n=1}^N a_n \cos nx - \sum_{n=1}^N b_n \sin nx\|_2 = 0$. If f is continuous with continuous derivative, then the convergence holds also pointwise.

We introduce the notion of weak derivative for measurable functions (see [1, Section 8.1]).

Definition 4.14. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a measurable function then f has weak derivative given by a measurable function w if and only if for every $\phi \in C_c^\infty(\mathbb{R})$, there holds

$$\int_{\mathbb{R}} f(x) \frac{d\phi(x)}{dx} dx = - \int_{\mathbb{R}} w(x) \phi(x) dx.$$

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a measurable function then f has weak gradient given by a measurable function $w = (w_1, \dots, w_d)$, $w_i : \mathbb{R}^d \rightarrow \mathbb{R}$, if and only if for every $\phi \in C_c^\infty(\mathbb{R})$, there holds

$$\int_{\mathbb{R}^d} f(x) \partial_i \phi(x) dx = - \int_{\mathbb{R}^d} w_i(x) \phi(x) dx.$$

It is possible to show that if the weak derivative (or weak gradient) exists, then it is unique. We will denote it as f' (or Df) to be intended in weak sense.

Example 4.15. Obviously if f is also differentiable in the usual sense, then $Df = \nabla f$ (it is just an application of the integration by parts formula).

Let now fix $f(x) = |x|$ for $x \in \mathbb{R}$ and compute the weak derivative. Let us fix a function ϕ which is smooth, has compact support. Then applying the definition we get, using the integration by part formula,

$$\begin{aligned} \int_{-\infty}^{+\infty} \phi(x)f'(x)dx &= - \int_{-\infty}^0 \phi'(x)(-x)dx - \int_0^{+\infty} \phi'(x)x dx \\ &= \int_{-\infty}^0 \phi'(x)x dx - \int_0^{+\infty} \phi'(x)x dx \\ &= [x\phi(x)]_{-\infty}^0 - \int_{-\infty}^0 \phi(x)dx - [x\phi(x)]_0^{+\infty} + \int_0^{+\infty} \phi(x)dx \\ &= - \int_{-\infty}^0 \phi(x)dx + \int_0^{+\infty} \phi(x)dx. \end{aligned}$$

Therefore, the weak derivative of f is the Heaviside function

$$f'(x) = H(x) = \begin{cases} 1 & x > 0 \\ -1 & x < 0. \end{cases}$$

We may wonder if also this function admits a weak derivative $H'(x)$. If this exists by definition there holds that for every smooth function with compact support

$$\int_{\mathbb{R}} H'(x)\phi(x)dx = - \int_{\mathbb{R}} \phi'(x)H(x)dx = + \int_{-\infty}^0 \phi'(x)dx - \int_0^{+\infty} \phi'(x)dx = \phi(0) + \phi(0) = 2\phi(0)$$

where in the second equality we used the definition of H and then we used the fact that $\phi = 0$ for $x \rightarrow +\infty, -\infty$.

Therefore

$$\int_{\mathbb{R}} H'(x)\phi(x)dx = 2\phi(0)$$

for every smooth function for compact support. This means that $H'(x)$ is an object which is zero everywhere except at zero, but its integral against a test function ϕ is nonzero. This contradicts our notion, that integrable functions can be changed on a set of measure zero without changing the integral. Indeed, H' is not a function in the usual sense, so **the Heaviside function has not weak derivative**. We may say that H has a derivative in the sense of distribution (so the derivative of H is a measure), and we write $H'(x) = 2\delta_0(x)$, where δ_0 is the Dirac δ -distribution, which associate to every smooth function its value at 0.

We introduce then the Sobolev space, for a given open set $O \subseteq \mathbb{R}^d$

$$H^1(O) = \{f \in L^2(O) \mid f \text{ admits a weak gradient } Df = (\partial_1 f, \dots, \partial_d f), \text{ with } w_i \in L^2(O) \text{ for all } i\}.$$

We put on H^1 the following scalar product: for $f, g \in H^1(O)$,

$$(f, g) = \int_O \left(f(x)g(x) + \sum_i \partial_i f(x)\partial_i g(x) \right) dx.$$

Note that it is well posed and it is a scalar product (since $\partial_i f, \partial_i g$ are in L^2 , then we may use the Holder inequality to show that $\partial_i f, \partial_i g \in L^1$).

The norm associated with this scalar product is

$$\left(\int_{\mathbb{R}^d} |f(x)|^2 + |Df(x)|^2 dx \right)^{1/2}.$$

it is easy to show that this norm is equivalent to the following one (which is the norm we are going to use:

$$\|f\|_{H^1} = \|f\|_{L^2} + \|Df\|_{L^2}.$$

It is possible to show that the space H^1 with this norm is a Banach space, so, we defined an Hilbert space (see [1, Thm 8.21]).

4.5 Bounded linear operators in Hilbert spaces

Let H be a Hilbert space. We consider linear bounded operators $T : H \rightarrow H$.

Definition 4.16 (Adjoint of an operator). *Let $T : H \rightarrow H$ be a bounded linear operator. The adjoint of T is the operator $T^* : H \rightarrow H$ such that $(Th, k) = (h, T^*k)$ for all $h, k \in H$. T is symmetric if $T = T^*$.*

Proposition 4.17. *Let T be a linear bounded symmetric operator. Then $\|T\| = \sup_{\|x\|=1} |(Tx, x)|$.*

Proof. By Cauchy Schwartz inequality we get

$$|(Tx, x)| \leq \|Tx\| \|x\| \leq \|T\| \|x\|^2 = \|T\| \|x\|^2.$$

On the other hand take $x \in H$ with $\|x\| = 1$ and $Tx \neq 0$ and define $y = Tx/\|Tx\|$. Then $\|y\| = 1$ and by symmetry and linearity of the operator

$$(Ty, x) = \frac{1}{\|Tx\|} (T(Tx), x) = \frac{1}{\|Tx\|} (Tx, Tx) = \|Tx\|.$$

A simple computation gives that $4(Ty, x) = (T(x+y), x+y) - (T(x-y), x-y)$, and then we get

$$\begin{aligned} 4\|Tx\| &= 4|(Ty, x) = (T(x+y), x+y) - (T(x-y), x-y) \leq \sup_{\|z\|=1} |(Tz, z)| (\|x+y\|^2 + \|x-y\|^2) \\ &= \sup_{\|z\|=1} |(Tz, z)| (2\|x\|^2 + 2\|y\|^2) \end{aligned}$$

where at the end we used the parallelogram identity. So we deduce, recalling that $\|x\| = 1 = \|y\|$ that $\|Tx\| \leq \sup_{\|z\|=1} |(Tz, z)|$. This gives the conclusion taking the supremum with respect to x . \square

Definition 4.18 (Compact operators). *Let $T : H \rightarrow H$ be a linear bounded operator. T is compact if for every bounded sequence $(h_n)_n$, there exists a subsequence such that $(Th_n)_n$ has a limit, that is $\lim_n Th_n = v$.*

Equivalently (it has to be proved though), an operator is compact if for every sequence $h_n \rightarrow h$ (h_n is weakly converging to h), there holds that $\lim_n \|Th_n - Th\| = 0$, so Th_n converge strongly to Th .

Definition 4.19 (Point spectrum (eigenvalues) of an operator). *The **point spectrum** $\sigma_p(T)$ of a operator is given by the **eigenvalues** of T , that is by the elements $\lambda \in \mathbb{R}$ such that there exists $v \in H$ (called **eigenvector**) for which $Tv = \lambda v$:*

$$\sigma_p(T) := \{\lambda \in \mathbb{R} \mid \exists v \in H, Tv = \lambda v\}.$$

*Given $\lambda \in \sigma_p(T)$, every element $v \in H$ such that $Tv = \lambda v$ is called **eigenvector** relative to the eigenvalue λ .*

*The **kernel** of an operator is the subspace N of H composed by vectors $h \in H$ such that $Th = 0$ (N is the space of eigenvectors relative to the eigenvalue 0).*

Theorem 4.20 (Spectral theorem for compact symmetric operators). *Let T be a symmetric compact operator on H , separable Hilbert space.*

If H is infinite dimensional, then T is not invertible (even if the kernel of T can be 0).

There exist $x_k \in H$, $\lambda_k \in \mathbb{R}$, such that $\{x_k\}$ is an orthonormal basis of H and $Tx_k = \lambda_k x_k$ (that is λ_k are eigenvalues of T with associated eigenvectors x_k) and the space $\{x \in H \mid Tx = \lambda_k x\}$ for every $\lambda_k \neq 0$ has finite dimension (so the multiplicity of every non zero eigenvalue is finite). Moreover the set of all the eigenvalues $\{\lambda_k\}$ of T is either finite or countable, and in this case $\lim_k \lambda_k = 0$:

$$\text{either } \sigma_p(T) = \{\lambda_1, \dots, \lambda_N\} \quad \text{or } \sigma_p(T) = \{\lambda_k, k \in \mathbb{N}\} \text{ and } \lim_{k \rightarrow +\infty} \lambda_k = 0.$$

Finally, let

$$m = \inf_{\{h \in H \mid \|h\|=1\}} (Th, h) \quad M = \sup_{\{h \in H \mid \|h\|=1\}} (Th, h).$$

Then $m, M \in \sigma_p(T)$ and $\sigma_p(T) \subseteq [m, M]$.

Proof. For the proof we refer to [1, Theorem 6.3, Lemma 6.5]. \square

Remark 4.21. Let T be a symmetric compact operator and $\{\lambda_k\}$ the set of all eigenvalues of T . Let $V_k = \{x \in H, Tx = \lambda_k x\}$ for every k , and P_k be the projection on V_k . Then

$$T = \sum_k \lambda_k P_k.$$

Definition 4.22 (Hilbert-Schmidt operator). Let H be a separable Hilbert space and T a compact symmetric operator. We say that T is an Hilbert Schmidt operator if

$$\sum_k \lambda_k^2 = \sum_k \|Tv_k\|^2 < +\infty$$

where λ_k are the eigenvalues of T .

Actually it can be proved that if T is a Hilbert Schmidt operator, the value of the sum $\sum_k \|Tv_k\|^2$ does not depend on the choice of the orthonormal basis v_k .

Proposition 4.23. Let H be a separable metric space with orthonormal basis u_i and \mathcal{H} be the space of all Hilbert Schmidt operators. Then this space with the norm

$$\|T\| = \sum_i \|Tu_i\|$$

is a Hilbert space with scalar product given by

$$(S, T) = \sum_i (Su_i, Tu_i).$$

Example 4.24. [Finite dimensional case] Let $H = \mathbb{R}^n$ and $Tx = \mathbf{A}x$ for some $n \times n$ matrix \mathbf{A} with values in \mathbb{R} . The adjoint of T is $T^*x = \mathbf{A}^T x$, where \mathbf{A}^T is the traspose of the matrix \mathbf{A} . T is symmetric if and only if \mathbf{A} is symmetric. Moreover the eigenvalues of T are the eigenvalues of the matrix \mathbf{A} .

Finally the spectral theorem for compact symmetric operators says that if \mathbf{A} is a symmetric matrix, then it can be reduced to diagonal form by a orthogonal transformation.

Remark 4.25. Let T be a Hilbert-Schmidt operator, such that 1 is not an eigenvalue of T . Then for all $f \in H$, the equation

$$h - Th = f$$

admits a unique solution $h \in H$. Indeed, consider v_k an orthonormal basis of H composed by eigenvectors of T . Then we rewrite the equation as

$$h - Th = \sum_k (h, v_k)v_k - \sum_k (h, v_k)\lambda_k v_k = \sum_k (1 - \lambda_k)(h, v_k)v_k = f = \sum_k (f, v_k)v_k.$$

Therefore the equation is satisfied if

$$(1 - \lambda_k)(h, v_k) = (f, v_k) \quad \text{that is } (h, v_k) = \frac{(f, v_k)}{1 - \lambda_k}.$$

Then the solution h to the equation is given by

$$h := \sum_k \frac{(f, v_k)}{1 - \lambda_k} v_k.$$

4.6 Problems

- (i) Let $X_n, Y_n \in \mathcal{H}$ such that $X_n \rightarrow X$ and $Y_n \rightarrow Y$. Show that
- $\mathbb{E}(X_n) \rightarrow \mathbb{E}(X)$,
 - $(X_n, Y_n) = \mathbb{E}(X_n Y_n) \rightarrow \mathbb{E}(XY) = (X, Y)$,
 - $Cov(X_n, Y_n) = \mathbb{E}(X_n Y_n) - \mathbb{E}(X_n)\mathbb{E}(Y_n) \rightarrow Cov(XY) = \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y)$
 - $Var(X_n) = Cov(X_n, X_n) \rightarrow Var(X) = Cov(X, X)$.
- (ii) Consider $H = L^2(-1, 1)$.
- (a) Let $V_1 = \{a + bx \mid a, b \in \mathbb{R}, x \in (-1, 1)\}$ (the subspace of polynomials of degree less than 1.) Find the orthogonal projection of x^2 on V_1 .
 - (b) Let $V_2 = \{a + bx + cx^2 \mid a, b, c \in \mathbb{R}, x \in (-1, 1)\}$ (the subspace of polynomials of degree less than 2). Find the orthogonal projection of x^3 on V_2 .
- (iii) Consider $X, Y, Z \in \mathcal{H}$ and assume X, Z are not constant. Compute the least linear quadratic estimator $L(Y|X, Z)$. Show that $L(Y|X, Z) = L(Y|X) + L(Y|Z - L(Z|X)) - \mathbb{E}(Y)$. (Hint: look at Remark 4.12).
- (iv) Let $T : L^2(0,1) \rightarrow L^2(0,1)$ defined as $Tf(x) = \int_0^x f(y)dy$. Show that this is a compact operator and compute its adjoint.

References

- [1] A. Bressan *Lecture notes on Functional Analysis, with applications to Linear Partial Differential Equations* Graduate Studies in Mathematics, vol 143, AMS, 2013.
- [2] G. Folland *Real Analysis: modern techniques and their applications*. Wiley 1999 (2nd ed).

A Solutions to problems Section 2

- (i) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a monotone function. Show that f is Lebesgue measurable.

It is sufficient to show that for all $c \in \mathbb{R}$ the set $\{x \in \mathbb{R} \mid f(x) > c\}$ is measurable.

Assume that f is monotone increasing (if it is monotone decreasing the argument is analogous). Let $c \in \mathbb{R}$. If $f(x) \leq c$ for all $x \in \mathbb{R}$ then $\{x \in \mathbb{R} \mid f(x) > c\}$ is the empty set and we are done.

Assume now that there exists $\bar{x} \in \mathbb{R}$ such that $f(\bar{x}) > c$. By monotonicity we get that $f(y) > c$ for all $y > \bar{x}$. We consider now the set $A_c = \{x \in \mathbb{R} \mid f(x) > c\}$. Our aim is to show that this is a measurable set.

We observed that by monotonicity, if $x \in A_c$, then $[x, +\infty) \subseteq A_c$. So, if A_c is not bounded from below, this implies that $A_c = \mathbb{R}$ and so we are done. Assume now that A_c is bounded from below and define $x_c = \inf A_c$. For all $x > x_c$ we get that $f(x) > c$ and $f(x) \leq c$ for all $x < x_c$. This implies that $A_c = (x_c, +\infty)$ if $f(x_c) \leq c$, and $A_c = [x_c, +\infty)$ if $f(x_c) > c$. In both cases, $A_c \in \mathcal{M}$.

Note that actually we get something more: for all c , we get that A_c is a Borel set, so the function f is Borel measurable.

- (ii) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a Lebesgue measurable function and let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $f(x) = g(x)$ for almost every $x \in \mathbb{R}^n$. Show that g is Lebesgue measurable.

Let $c \in \mathbb{R}$ and define $A_c = \{x \mid f(x) > c\}$ and $B_c = \{x \mid g(x) > c\}$. Then, by assumption $(A_c \setminus B_c) \cup (B_c \setminus A_c)$ has measure 0. This implies that if $A_c \in \overline{\mathcal{M}}$ for all c , then $B_c \in \overline{\mathcal{M}}$ for all c , and so g is measurable.

- (iii) Consider the right continuous increasing function on \mathbb{R}

$$F(x) = \begin{cases} x & x < 0 \\ x + 1 & x \geq 0. \end{cases}$$

Which is the Borel measure associated to this function?

We define $\mu_F(a, b] = F(b) - F(a)$, and then we extend it to a measure on the Borel σ -algebra. Given F as in the statement, we get that $\mu_F(a, b] = b - a$ if $a < b < 0$, $\mu_F(a, b] = b + 1 - (a + 1) = b - a$ if $0 \leq a < b$, whereas if $a < 0 \leq b$, then $\mu_F(a, b] = b + 1 - a = b - a + 1$. Therefore $\mu_F = \mathcal{L} + \delta_0$.

- (iv) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a positive function. Let $F(t)$ the repartition function of f . Recall that $\int_{\mathbb{R}^n} f(x) dx = \int_0^{+\infty} F(t) dt$. Show that for all $p > 1$

$$\int_{\mathbb{R}^n} |f(x)|^p dx = p \int_0^{+\infty} t^{p-1} F(t) dt.$$

So $f \in L^p(\mathbb{R}^n)$ if and only if $p \int_0^{+\infty} t^{p-1} F(t) dt < +\infty$.

We know that $\int_{\mathbb{R}^n} |f(x)|^p dx = \int_0^{+\infty} F_p(t) dt$, where F_p is the repartition function of $|f|^p$, that is

$$F_p(t) = \mathcal{L}\{x \mid |f(x)|^p > t\} = \mathcal{L}\{x \mid |f(x)| > t^{1/p}\} = F(t^{1/p}).$$

Therefore

$$\int_{\mathbb{R}^n} |f(x)|^p dx = \int_0^{+\infty} F_p(t) dt = \int_0^{+\infty} F(t^{1/p}) dt = \int_0^{+\infty} F(s) p s^{p-1} ds$$

where the last equality comes from the change of variable $s^p = t$.

(v) Let

$$f(x) = \begin{cases} \frac{1}{|x|} & |x| < 1 \\ 0 & |x| > 1. \end{cases}$$

For which p , is the function $f \in L^p(\mathbb{R}^n)$? For such p , compute $\|f\|_p$.

Observe that

$$A(t) = \{x \mid f(x) > t\} = \begin{cases} \{x \mid |x| < t^{-1}\} & t > 1 \\ \{x \mid |x| < 1\} & 0 < t \leq 1 \end{cases}$$

So in \mathbb{R}^n , we get that the repartition function is given by

$$F(t) := \mathcal{L}(A(t)) = \begin{cases} \mathcal{L}(B(0, t^{-1})) = \mathcal{L}(B(0, 1)) t^{-n} & t > 1 \\ \mathcal{L}(B(0, 1)) & 0 < t \leq 1 \end{cases}.$$

Therefore by the previous exercise

$$\begin{aligned} \int_{\mathbb{R}^n} |f(x)|^p dx &= p \int_0^{+\infty} t^{p-1} F(t) dt = p \mathcal{L}(B(0, 1)) \left(\int_0^1 t^{p-1} dt + \int_1^{+\infty} t^{p-1-n} dt \right) = \\ &= \begin{cases} p \mathcal{L}(B(0, 1)) \left(\frac{1}{p} + \frac{1}{n-p} \right) = \frac{n}{n-p} \mathcal{L}(B(0, 1)) & p - n < 0 \\ +\infty & p - n \geq 0 \end{cases}. \end{aligned}$$

Therefore $f \in L^p(\mathbb{R}^n)$ if and only if $p < n$ and $\|f\|_p = \left(\frac{n}{n-p} \mathcal{L}(B(0, 1)) \right)^{1/p}$.

Finally we observe that $f \notin L^\infty(\mathbb{R}^n)$.

(vi) Let

$$f(x) = \begin{cases} \log\left(\frac{1}{|x|}\right) & 0 < |x| < 1 \\ 0 & |x| = 0. \end{cases}$$

Show that $f \in L^1(B(0, 1))$.

Is it true that the function f satisfies $f \in L^p(B(0, 1))$ for every $p \geq 1$?

Is it true that $f \in L^\infty(B(0, 1))$?

Observe that $f \geq 0$ and for $t > 0$,

$$A(t) = \{x \mid f(x) > t\} = \{x \mid |x| < e^{-t}\}.$$

So in \mathbb{R}^n , we get

$$\mathcal{L}(A(t)) = \mathcal{L}(B(0, 1)) e^{-tn}.$$

Applying the definition we get

$$\int_{\mathbb{R}^n} f(x) dx = \int_0^{+\infty} \mathcal{L}(A(t)) dt = \mathcal{L}(B(0, 1)) \int_0^{+\infty} e^{-tn} dt = \frac{\mathcal{L}(B(0, 1))}{n} < \infty.$$

So $f \in L^1(\mathbb{R}^n)$.

Analogously it is possible to show that $f \in L^p(\mathbb{R}^n)$ since

$$\int_{\mathbb{R}^n} |f(x)|^p dx = p \mathcal{L}(B(0, 1)) \int_0^{+\infty} t^{p-1} e^{-tn} dt < \infty.$$

Finally observe that $f \notin L^\infty(\mathbb{R}^n)$ since $\lim_{|x| \rightarrow 0} f(x) = +\infty$.

(vii) Let

$$f(x) = \begin{cases} \frac{1}{|x|} & |x| > 1 \\ 1 & |x| \leq 1. \end{cases}$$

For which p , is the function f in the space $L^p(\mathbb{R}^n)$?

According to the definition of $L^p(\mathbb{R}^n)$, we have to check for which p , the function $|f|^p$ is summable in \mathbb{R}^n .

Observe that

$$A(t) = \{x \mid |f(x)|^p > t\} = \begin{cases} \emptyset & t \geq 1 \\ \{x \mid |x| < t^{-\frac{1}{p}}\} & t < 1. \end{cases}$$

So

$$\mathcal{L}(A(t)) = \begin{cases} 0 & t \geq 1 \\ \mathcal{L}(B(0, 1))t^{-\frac{n}{p}} & t < 1. \end{cases}$$

Applying the definition we get

$$\begin{aligned} \int_{\mathbb{R}^n} |f(x)|^p dx &= \int_0^{+\infty} \mathcal{L}(A(t)) dt = \int_0^1 \mathcal{L}(B(0, 1)) t^{-\frac{n}{p}} dt = \mathcal{L}(B(0, 1)) \left. \frac{1}{1 - \frac{n}{p}} t^{1 - \frac{n}{p}} \right|_0^1 \\ &= \begin{cases} +\infty & p \leq n \\ \mathcal{L}(B(0, 1)) \frac{p}{p-n} & p > n. \end{cases} \end{aligned}$$

So the function $f \in L^p(\mathbb{R}^n)$ for $p > n$. Note that $f \in L^\infty(\mathbb{R}^n)$ and $\|f\|_\infty = 1$.

B Solutions to problems Section 3

(i) Let $(X, \|\cdot\|)$ a Banach space and $F : X \rightarrow X$ such that there exists $0 < a < 1$ for which

$$\|F(x) - F(y)\| \leq a\|x - y\| \quad \forall x, y \in X.$$

(F is a contraction)

(a) Show that the map F is continuous.

(b) Let $x_0 \in X$. Define $x_1 = F(x_0)$, $x_2 = F(x_1)$ and so on $x_n = F(x_{n-1})$. Prove that

$$\|x_n - x_{n+1}\| \leq a^n \|x_0 - x_1\|.$$

Deduce that $(x_n)_n$ is a Cauchy sequence.

(c) Let $\bar{x} = \lim_n x_n$, where (x_n) has been defined in the previous step. Show that $F(\bar{x}) = \bar{x}$. So, \bar{x} is a fixed point of F .

(d) Show that the map F admits a unique fixed point, that is a point such that $\bar{x} = F(\bar{x})$.

This is called **Banach-Caccioppoli theorem**.

(a) Let (x_n) be a sequence in X which is converging to x . Then $0 \leq \|F(x_n) - F(x)\| \leq a\|x_n - x\|$, and so $\lim_{n \rightarrow +\infty} F(x_n) = F(x)$ since $\lim_{n \rightarrow +\infty} x_n = x$.

(b) By the property of the function F and the definition of the we get that

$$\begin{aligned} \|x_{n+1} - x_n\| &= \|F(x_n) - F(x_{n-1})\| \leq a\|x_n - x_{n-1}\| = \\ &= a\|F(x_{n-1}) - F(x_{n-2})\| \leq a^2\|x_{n-1} - x_{n-2}\| \leq \dots \leq a^n\|x_1 - x_0\|. \end{aligned}$$

Let $n > m$. Then, by using the triangular inequality, we get

$$\|x_n - x_m\| \leq \|x_n - x_{n-1}\| + \|x_{n-1} - x_{n-2}\| + \cdots + \|x_{m+1} - x_m\|.$$

By using the previous inequality and recalling that $\sum_{i=0}^n a^i = \frac{1-a^{n+1}}{1-a}$, we get

$$\|x_n - x_m\| \leq (a^n + a^{n-1} + \cdots + a^m) \|x_0 - x_1\| \leq \frac{a^{m+1} - a^{n+1}}{1-a} \|x_0 - x_1\|.$$

Since $0 < a < 1$, we get that $a^{n+1}, a^{m+1} \rightarrow 0$ as $n, m \rightarrow +\infty$. So in particular the previous inequality implies that (x_n) is a Cauchy sequence.

- (c) Since (x_n) is a Cauchy sequence, and the space is complete, it is converging to some point x . Using the continuity of F we have that $\lim_n F(x_n) = F(x)$. But we recall that $\lim_n F(x_n) = \lim_n x_{n-1} = x$. So $F(x) = x$.
- (d) Let x, z such that $F(x) = x$ and $F(z) = z$. Then by the property of F , and recalling that $a < 1$,

$$\|x - z\| = \|F(x) - F(z)\| \leq a\|x - z\| < \|x - z\|.$$

This is not possible unless $\|x - z\| = 0$, which implies $z = x$.

- (ii) Let $f \in L^p(\mathbb{R}^n)$ and $\alpha > 0$. Prove that

$$\mathcal{L}(\{x \in \mathbb{R}^n \mid |f(x)| > \alpha\})^{\frac{1}{p}} \leq \frac{1}{\alpha} \|f\|_p.$$

This is called **Chebyshev inequality**.

Let $A_\alpha = \{x \in \mathbb{R}^n \mid |f(x)| > \alpha\}$. Then $\mathbb{R}^n = A_\alpha \cup (\mathbb{R}^n \setminus A_\alpha)$. So we compute, recalling definitions,

$$\|f\|_p^p = \int_{\mathbb{R}^n} |f(x)|^p dx = \int_{A_\alpha} |f(x)|^p dx + \int_{\mathbb{R}^n \setminus A_\alpha} |f(x)|^p dx \geq \int_{A_\alpha} |f(x)|^p dx,$$

since $|f|^p \geq 0$. Now if $x \in A_\alpha$, then $|f(x)|^p \geq \alpha^p$. Therefore in the previous inequality we get

$$\|f\|_p^p \geq \int_{A_\alpha} |f(x)|^p dx \geq \alpha^p \int_{A_\alpha} dx = \alpha^p \mathcal{L}(A_\alpha).$$

This gives the desired inequality,

$$\mathcal{L}(A_\alpha) \leq \left(\frac{\|f\|_p}{\alpha} \right)^p$$

after extracting the p root.

- (iii) Let $f \in L^\infty(O) \cap L^p(O)$ for some $p > 1$.

- (a) Show that $f \in L^q(O)$ for every $q \geq p$.
- (b) Prove that $\lim_{q \rightarrow +\infty} \|f\|_q \leq \|f\|_\infty$.
- (c) By using the Chebyshev inequality prove that for every $a > 0$ and every $q \geq p$

$$a \mathcal{L}(\{x \in \mathbb{R}^n \mid |f(x)| > a\})^{\frac{1}{q}} \leq \|f\|_q.$$

Observe that if $\mathcal{L}(\{x \in \mathbb{R}^n \mid |f(x)| > a\}) \neq 0$, then $a \leq \lim_{q \rightarrow +\infty} \|f\|_q$.

- (d) Using the previous point show that $\lim_{q \rightarrow +\infty} \|f\|_q \geq \|f\|_\infty$ and therefore by point 2, $\lim_{q \rightarrow +\infty} \|f\|_q = \|f\|_\infty$

- (a) Since $f \in L^\infty$, we get that $|f(x)| \leq \|f\|_\infty$ for almost every x . Take $q > p$ and observe that $|f(x)|^q = |f(x)|^p |f(x)|^{q-p} \leq |f(x)|^p \|f\|_\infty^{q-p}$ for a.e. x . This implies that

$$\int_O |f(x)|^q dx \leq \int_O |f(x)|^p \|f\|_\infty^{q-p} dx = \|f\|_\infty^{q-p} \int_O |f(x)|^p dx < \infty$$

and so $f \in L^q(\mathbb{R}^n)$.

- (b) By the previous inequality we get

$$\|f\|_q^q = \int_O |f(x)|^q dx \leq \|f\|_\infty^{q-p} \|f\|_p^p.$$

We extract the q -root of both terms and we get

$$\|f\|_q \leq \|f\|_\infty^{1-\frac{p}{q}} \|f\|_p^{\frac{p}{q}}.$$

So taking the limit as $q \rightarrow +\infty$ (recalling the p is fixed!) we get $\lim_{q \rightarrow +\infty} \|f\|_q \leq \|f\|_\infty$.

- (c) It is just Chebycheff inequality. If $\mathcal{L}(\{x \in \mathbb{R}^n \mid |f(x)| > a\}) \neq 0$, then sending $q \rightarrow +\infty$, we conclude.
- (d) By previous inequality we get

$$\|f\|_q \geq (\|f\|_\infty - \varepsilon) |A_\varepsilon|^{\frac{1}{q}},$$

where $A_\varepsilon = \{x \in \mathbb{R}^n \mid |f(x)| > \|f\|_\infty - \varepsilon\}$. Now, by definition of $\|f\|_\infty$, we get that for every $\varepsilon > 0$, $\mathcal{L}(A_\varepsilon) > 0$ (otherwise if $\mathcal{L}(A_\varepsilon) = 0$, then the right L^∞ norm should be $\|f\|_\infty - \varepsilon$.) This implies that for every $\varepsilon > 0$ fixed $\lim_{q \rightarrow +\infty} \mathcal{L}(A_\varepsilon)^{\frac{1}{q}} = 1$. Taking the limit in the previous inequality then we have

$$\lim_{q \rightarrow +\infty} \|f\|_q \geq \|f\|_\infty - \varepsilon.$$

Now taking the limit $\varepsilon \rightarrow 0$, we obtain the desired inequality $\lim_{q \rightarrow +\infty} \|f\|_q \geq \|f\|_\infty$, which, together with 2 gives the conclusion.

- (iv) Prove that if $f \in L^2(-1, 1)$ then $f \in L^1(-1, 1)$ and moreover

$$\|f\|_1 \leq \sqrt{2} \|f\|_2.$$

Provide an example of a function $f \in L^1(-1, 1)$ such that $f \notin L^2(-1, 1)$.

Since $\chi_{(-1,1)} \in L^2(-1, 1)$, then by Holder inequality $f = f\chi_{(-1,1)} \in L^1(-1, 1)$. Moreover

$$\|f\|_1 \leq \|f\|_2 \left(\int_{-1}^1 dx \right)^{\frac{1}{2}} = \sqrt{2} \|f\|_2.$$

The function $f(x) = \frac{1}{\sqrt{|x|}}$ is in $L^1(-1, 1)$ but not in $L^2(-1, 1)$.

- (v) Let $f_n \in L^p(O)$ such that $f_n \rightharpoonup f$ weakly in L^p for $p > 1$. Show that there exists $C > 0$ such that $\|f_n\|_p \leq C$.

Let q be the adjoint of p . Define the operator $T_n : L^q(O) \rightarrow \mathbb{R}$ defined as $T_n g = \int_O f_n(x) g(x) dx$ and $Tg = \int_O f(x) g(x) dx$. For all n , by Holder inequality this is a linear bounded operator with norm $\|T_n\| = \|f_n\|_p$. Observe that since $f_n \rightharpoonup f$, we have that for all $g \in L^q$, there holds $T_n g \rightarrow Tg$, so in particular for all $g \in L^q$ there exists $C_g > 0$ such that $\|T_n g\| \leq C_g$. We conclude by the uniform boundedness principle.

(vi) Consider the following operator $T : L^2(0, 2) \rightarrow L^2(0, 2)$ defined as

$$Tf(x) = \int_0^x f(y)dy.$$

Show that this is a bounded continuous operator.

First of all T is linear by linearity of the integral.

Note that, by Jensen, and changing the order of integration (observe that $0 < y < x < 2$) we get

$$\begin{aligned} \|Tf\|_2^2 &= \int_0^2 \left(\int_0^x \chi_{(0,x)}(y)f(y)dy \right)^2 dx \leq \int_0^2 2 \int_0^x \chi_{(0,x)}(y)|f(y)|^2 dy = 2 \int_0^2 \int_0^x |f(y)|^2 dy \\ &= 2 \int_0^2 \int_y^2 |f(y)|^2 dx dy = 2 \int_0^2 (2-y)|f(y)|^2 dy \leq 4\|f\|_2^2 \end{aligned}$$

where the last inequality comes from the fact that $(2-y) \leq 2$ for $y \in (0, 2)$. Therefore

$$\sup_{\{\|f\| \leq 1\}} \|Tf\|_2 \leq \sup_{\{\|f\| \leq 1\}} 2\|f\|_2 \leq 2.$$

(vii) Let $O \subseteq \mathbb{R}^n$ be a open set, $f_k, f \in L^p(O)$, for all $k \in \mathbb{N}$.

(a) Show that if $f_k \rightarrow f$ (strongly) in $L^p(O)$ then $\lim_k \|f_k\|_p = \|f\|_p$.

(b) Show that if $f_k \rightharpoonup f$ (weakly) in $L^p(O)$ then $\liminf_k \|f_k\|_p \geq \|f\|_p$.

(a) By triangle inequality we get that

$$-\|f_k - f\|_p \leq \|f_k\|_p - \|f\|_p \leq \|f_k - f\|_p$$

from which we conclude passing to the limit.

(b) We define the function

$$\text{sign}f(x) = \begin{cases} \frac{f(x)}{|f(x)|} & f(x) \neq 0 \\ 0 & f(x) = 0, \end{cases}$$

Note that $f(x)\text{sign}f(x) = |f(x)|$. Let $g(x) = |f(x)|^{p-1}\text{sign}f(x)$. Then $g \in L^q(O)$ for $q = \frac{p}{p-1}$, and $\|g\|_q = \|f\|_p^{\frac{q}{p}}$. Finally note that $g(x)f(x) = |f(x)|^p$. Then, by weak convergence and Holder inequality, we get that

$$\begin{aligned} \|f\|_p &= \int_O g(x)f(x)dx = \lim_k \int_O f_k(x)g(x)dx \\ &\leq \liminf_k \|f_k\|_p \|g\|_q = \liminf_k \|f_k\|_p \|f\|_p^{\frac{q}{p}} \end{aligned}$$

which gives the conclusion.

(viii) Let $O \subseteq \mathbb{R}^n$ be a open set, let $p \geq 1$ and $q = \frac{p}{p-1}$, its conjugate exponent.

Let $f_k, f \in L^p(O)$, for all $k \in \mathbb{N}$, $g_k, g \in L^q(O)$, for all $k \in \mathbb{N}$.

(a) Assume $f_k \rightarrow f$ (strongly) in $L^p(O)$ and $g_k \rightarrow g$ (strongly) in $L^q(O)$.

Show that $f_k g_k \rightarrow f g$ (strongly) in $L^1(O)$.

(b) Assume $f_k \rightarrow f$ (strongly) in $L^p(O)$ and $g_k \rightharpoonup g$ (weakly) in $L^q(O)$ (this implies in particular that $\|g_k\|_q \leq M$).

Show that $f_k g_k \rightarrow f g$ (strongly) in $L^1(O)$.

(a) We have to prove that $\int_O |f_k(x)g_k(x) - f(x)g(x)|dx \rightarrow 0$ as $k \rightarrow +\infty$.

We observe that, by Holder inequality,

$$\begin{aligned} \int_O |f_k(x)g_k(x) - f(x)g(x)|dx &= \int_O |f(x)(g_k(x) - g(x)) + (f_k(x) - f(x))g_k(x)|dx \\ &\leq \int_O |f(x)||g_k(x) - g(x)|dx + \int_O |f_k(x) - f(x)||g_k(x)|dx \\ &\leq \|g_k - g\|_q \|f\|_p + \|f_k - f\|_p \|g_k\|_q. \quad (\text{B.1}) \end{aligned}$$

Sending $k \rightarrow +\infty$, and recalling that $\lim_k \|g_k\|_q = \|g\|_q$, we get the conclusion.

(b) Arguing as in (B.1), we get

$$\begin{aligned} \int_O |f_k(x)g_k(x) - f(x)g(x)|dx &\leq \int_O |f(x)||g_k(x) - g(x)|dx + \int_O |f_k(x) - f(x)||g_k(x)|dx \\ &\leq \int_O |f(x)||g_k(x) - g(x)|dx + \|f_k - f\|_p \|g_k\|_q. \end{aligned}$$

Note that $\int_O |f(x)||g_k(x) - g(x)|dx \rightarrow 0$ by weak convergence of g_k to g , whereas $\|f_k - f\|_p \rightarrow 0$ by strong convergence of f_k to f . Moreover recall that there exists $M > 0$ such that $\|g_k\|_q \leq M$. So, we conclude, sending $k \rightarrow +\infty$ that $\int_O |f_k(x)g_k(x) - f(x)g(x)|dx \rightarrow 0$.

(ix) Let $O \subseteq \mathbb{R}^n$ be a open set, and $f_k, f \in L^2(O)$, for all $k \in \mathbb{N}$ such that $f_k \rightharpoonup f$ weakly in $L^2(O)$.

Show that if $\lim_k \|f_k\|_2 = \|f\|_2$, then $f_k \rightarrow f$ strongly in $L^2(O)$.

We compute

$$\begin{aligned} \|f_k - f\|_2^2 &= \int_O |f_k(x) - f(x)|^2 dx = \int_O |f_k|^2 - 2 \int_O f_k(x)f(x)dx + \int_O |f(x)|^2 dx \\ &= \|f_k\|_2^2 - 2 \int_O f_k(x)f(x)dx + \|f\|_2^2. \end{aligned}$$

Note that by weak convergence, since $f \in L^2(O)$ we get that $\lim_k \int_O f_k(x)f(x)dx = \int_O |f(x)|^2 dx = \|f\|_2^2$ and by assumption $\lim_k \|f_k\|_2^2 = \|f\|_2^2$. Therefore, passing to the limit above we get

$$\lim_k \|f_k - f\|_2^2 = \|f\|_2^2 - 2\|f\|_2^2 + \|f\|_2^2 = 0.$$

C Solutions to problems Section 4

- (i)
- $\mathbb{E}(X_n) = (X_n, 1) \rightarrow (X, 1) = \mathbb{E}(X)$, by continuity of the scalar product (as a consequence of Cauchy Schwartz inequality).
 - $(X_n, Y_n) = (X_n - X, Y_n - Y) + (X, Y_n) + (X_n, Y) - (X, Y)$. We conclude observing that $(X_n - X, Y_n - Y) \rightarrow 0$, $(X_n, Y) \rightarrow (X, Y)$ and $(X, Y_n) \rightarrow (X, Y)$.
 - the convergence of covariance and variance are immediate consequences of the first two items.
- (ii) Consider $H = L^2(-1, 1)$.
- (a) Let $V_1 = \{a + bx \mid a, b \in \mathbb{R}, x \in (-1, 1)\}$ (the subspace of polynomials of degree less than 1.) Find the orthogonal projection of x^2 on V_1 .
- (b) Let $V_2 = \{a + bx + cx^2 \mid a, b, c \in \mathbb{R}, x \in (-1, 1)\}$ (the subspace of polynomials of degree less than 2). Find the orthogonal projection of x^3 on V_2 .

- (a) We look for an orthonormal basis of V_1 . A basis of V_1 is given by $1, x$. Note that $(x, 1) = \int_{-1}^1 x dx = 0$, so 1 and x are orthogonal. We have to normalize them. We compute $\int_{-1}^1 dx = 2$ and $\int_{-1}^1 |x|^2 dx = \frac{2}{3}$. Therefore an orthonormal basis of V_1 is given by $(\frac{1}{\sqrt{2}}, \frac{\sqrt{3}x}{\sqrt{2}})$. By the theorem on orthogonal projection we have that

$$P_{V_1}(x^2) = a_0 \frac{1}{\sqrt{2}} + a_1 \frac{\sqrt{3}x}{\sqrt{2}}$$

where $a_0 = (x^2, \frac{1}{\sqrt{2}})$ and $a_1 = (x^2, \frac{\sqrt{3}x}{\sqrt{2}})$. We compute

$$a_0 = \left(x^2, \frac{1}{\sqrt{2}}\right) = \frac{1}{\sqrt{2}} \int_{-1}^1 x^2 dx = \frac{2}{3\sqrt{2}} \quad a_1 = \left(x^2, \frac{\sqrt{3}x}{\sqrt{2}}\right) = \frac{\sqrt{3}}{\sqrt{2}} \int_{-1}^1 x^3 dx = 0.$$

Therefore

$$P_{V_1}(x^2) = \frac{2}{3\sqrt{2}} \frac{1}{\sqrt{2}} = \frac{1}{3}.$$

Another way to compute the orthogonal projection is just using the definition: we have to find the point in V_1 with minimal distance from x^2 . Every point in V_1 is defined as $a + bx$ for some a, b , so we have to solve the minimization problem

$$\begin{aligned} \min_{a,b \in \mathbb{R}} \|x^2 - a - bx\|^2 &= \min_{a,b} \int_{-1}^1 |x^2 - a - bx|^2 dx \\ &= \min_{a,b} \int_{-1}^1 (x^4 + a^2 + b^2 x^2 - 2ax^2 - 2bx^3 + 2abx) dx = \min_{a,b} \left(\frac{1}{5} + 2a^2 + \frac{b^2}{3} - \frac{2}{3}a \right). \end{aligned}$$

The minimum is $1/5$ and there are two minimum couples of points: $a = \frac{1}{3}, b = 0$ and $a = 0, b = 0$. Therefore the projection is $1/3 + 0x = 1/3$.

- (b) The previous point implies that $x^2 - \frac{1}{3} \in V_1^\perp$. We have to find a orthonormal basis of V_2 . In order to have three generators which are orthogonal, we consider $1, x, x^2 - \frac{1}{3}$. Moreover we normalize them to have norm 1. We compute

$$\int_{-1}^1 (x^2 - \frac{1}{3})^2 dx = \frac{2}{5} + \frac{2}{9} - \frac{4}{9} = \frac{2}{5} - \frac{2}{9} = \frac{8}{45}.$$

So an orthonormal basis of V_2 is given by $(\frac{1}{\sqrt{2}}, \frac{\sqrt{3}x}{\sqrt{2}}, \frac{\sqrt{45}}{\sqrt{8}}(x^2 - \frac{1}{3}))$. Again by theorem on orthogonal projections we have

$$P_{V_2}(x^3) = a_0 \frac{1}{\sqrt{2}} + a_1 \frac{\sqrt{3}x}{\sqrt{2}} + a_2 \frac{\sqrt{45}}{\sqrt{8}}(x^2 - \frac{1}{3}) = (t^3, \frac{\sqrt{3}t}{\sqrt{2}}) \frac{\sqrt{3}x}{\sqrt{2}}$$

where $a_0 = (x^3, \frac{1}{\sqrt{2}})$, $a_1 = (x^3, \frac{\sqrt{3}x}{\sqrt{2}})$, $a_2 = (x^3, \frac{\sqrt{45}}{\sqrt{8}}(x^2 - \frac{1}{3}))$. It is easy to check that $a_0 = 0 = a_2$. We compute

$$a_1 = (x^3, \frac{\sqrt{3}x}{\sqrt{2}}) = \frac{\sqrt{3}}{\sqrt{2}} \int_{-1}^1 x^4 dt = \frac{\sqrt{3}}{\sqrt{2}} \frac{2}{5}.$$

Therefore

$$P_{V_2}(x^3) = \frac{\sqrt{3}}{\sqrt{2}} \frac{2}{5} \frac{\sqrt{3}x}{\sqrt{2}} = \frac{3}{5}x.$$

(iii) Recalling Remark 4.12 we have that

$$L(Y|X, Z) = Pr_S(Y) = a + bX + cZ$$

where S is the space with basis $1, X, Z$.

Observe that by the same argument $L(Z|X) = Pr_T(Z)$ where T is the space with a basis given by $1, X$. In particular by Theorem 4.7 we have that $Z - L(Z|X) \in T^\perp$ and arguing as in Remark 4.12 $L(Z|X) = \mathbb{E}(Z) + \frac{Cov(X, Z)}{Var(X)}(X - \mathbb{E}(X))$.

An orthonormal basis of S can be therefore obtained by considering an orthonormal basis of T , which is given by $1, \frac{X - \mathbb{E}(X)}{\sqrt{Var(X)}}$ as proved in Remark 4.12 and then adding the element $k(Z - L(Z|X))$ where k is such that $\mathbb{E}(k(Z - L(Z|X)))^2 = 1$. Since $\mathbb{E}((Z - L(Z|X))^2) = \frac{Var(Z)Var(X) - Cov^2(X, Z)}{Var(X)}$ as proved in Remark 4.12, we get that $k = \frac{\sqrt{Var(X)}}{\sqrt{Var(Z)Var(X) - Cov^2(X, Z)}}$.

So, as in Remark 4.12,

$$\begin{aligned} L(Y|X, Z) &= \mathbb{E}(Y) + \frac{Cov(X, Y)}{Var(X)}(X - \mathbb{E}(X)) \\ &\quad + \frac{Var(X)Cov(Z, Y) - Cov(X, Z)Cov(X, Y)}{Var(Z)Var(X) - Cov^2(X, Z)}(Z - L(Z|X)) \\ &= \mathbb{E}(Y) \\ &\quad + \frac{Var(Z)Cov(X, Y) - Cov(Z, Y)Cov(X, Z)}{Var(Z)Var(X) - Cov^2(X, Z)}(X - \mathbb{E}(X)) \\ &\quad + \frac{Var(X)Cov(Z, Y) - Cov(X, Z)Cov(X, Y)}{Var(Z)Var(X) - Cov^2(X, Z)}(Z - \mathbb{E}(Z)). \end{aligned}$$

Observe that

$$\mathbb{E}(Y) + \frac{Cov(X, Y)}{Var(X)}(X - \mathbb{E}(X)) = L(Y|X)$$

and moreover

$$\mathbb{E}(Y) + \frac{Var(X)Cov(Z, Y) - Cov(X, Z)Cov(X, Y)}{Var(Z)Var(X) - Cov^2(X, Z)}(Z - L(Z|X)) = L(Y|Z - L(Z|X)).$$

This conclude the proof.

(iv) Let $T : L^2(0,1) \rightarrow L^2(0,1)$ defined as $Tf(x) = \int_0^x f(y)dy$.

Show that this is a compact operator and compute its adjoint.

To prove that it is compact we have to show that if f_n converge weakly to f , that is $\lim_n \int_0^1 f_n(x)g(x)dx = \int_0^1 f(x)g(x)dx$ for all g , then $\|Tf_n - Tf\|_2 \rightarrow 0$. We compute, reasoning as above,

$$\|Tf_n - Tf\|_2^2 = \int_0^1 \left(\int_0^1 \chi_{(0,x)}(y)(f_n(y) - f(y))dy \right)^2 dx.$$

Now $F_n(x) = \left(\int_0^1 \chi_{(0,x)}(y)(f_n(y) - f(y))dy \right)^2$ is a function such that $\lim_n F_n(x) = 0$ for all x (by weak convergence, since $\chi_{(0,x)} \in L^2$). Moreover by Jensen inequality $F_n(x) \leq \int_0^x |f_n(y) - f(y)|^2 dy \leq \|f_n - f\|_2^2 \leq (\|f_n\|_2 + \|f\|_2)^2$. Recall that since f_n converge weakly then there exists C such that $\|f_n\|_2 \leq C$ (see Problem v)). This implies that $0 \leq F_n(x) \leq 2C^2$. Since constant functions are element in $L^2(0,1)$, we conclude by Lebesgue dominated convergence that $\lim_n \|Tf_n - Tf\|_2^2 = 0$.

To compute the adjoint we recall that $(Tf, g) = (f, T^*g)$ and so we compute, changing the order of integration

$$(Tf, g) = \int_0^1 \int_0^x f(y)g(x)dydx = \int_0^1 \int_y^1 g(x)dx f(y)dy.$$

Therefore $T^*g(x) = \int_x^1 g(y)dy$.